

# Non-Noetherian generalized Heisenberg algebras

Samuel A. Lopes\*

## Abstract

In this note we classify the non-Noetherian generalized Heisenberg algebras  $\mathcal{H}(f)$  introduced in [8]. In case  $\deg f > 1$ , we determine all locally finite and also all locally nilpotent derivations of  $\mathcal{H}(f)$  and describe the automorphism group of these algebras.

**MSC Numbers (2010):** Primary 16W25, 16W20; Secondary 16P40, 16S36.

**Keywords:** generalized Heisenberg algebra; ambiskew polynomial ring; weak generalized Weyl algebra; automorphism group; derivation; Noetherian.

## 1 Introduction

Fix a polynomial  $f \in \mathbb{C}[h]$ . The *generalized Heisenberg algebra*  $\mathcal{H}(f)$  is the unital associative  $\mathbb{C}$ -algebra with generators  $x, y, h$  satisfying the relations:

$$hx = xf(h), \quad yh = f(h)y, \quad yx - xy = f(h) - h. \quad (1.1)$$

See [8] and the references therein for information on how these algebras first appeared and on their applications to theoretical physics.

Ambiskew polynomial rings were introduced by Jordan over a series of papers (see the references in [5]), but for our purposes the best suited definition is the one found in [5], which we briefly recall. Let  $\sigma$  be an endomorphism of a commutative  $\mathbb{C}$ -algebra  $B$ ,  $c \in B$  and  $p \in \mathbb{C}$ . The ambiskew polynomial ring  $R(B, \sigma, c, p)$  is the  $\mathbb{C}$ -algebra generated by  $B$  and two indeterminates,  $x$  and  $y$ , subject to the relations

$$bx = x\sigma(b), \quad yb = \sigma(b)y, \quad yx - pxy = c, \quad \text{for all } b \in B.$$

On comparing these relations with those in (1.1), one immediately sees that

$$\mathcal{H}(f) \cong R(\mathbb{C}[h], \sigma, f(h) - h, 1), \quad (1.2)$$

where  $\sigma : \mathbb{C}[h] \rightarrow \mathbb{C}[h]$  is the algebra endomorphism given by  $\sigma(h) = f(h)$ . In particular, one can see that there is an overlap between the generalized Heisenberg algebras defined above and (generalized) down-up algebras (see Corollary 2.7 below).

---

\*The author was partially supported by CMUP (UID/MAT/00144/2013), which is funded by FCT (Portugal) with national (MEC) and European structural funds through the programs FEDER, under the partnership agreement PT2020.

The algebras  $\mathcal{H}(f)$  can also be seen as weak generalized Weyl algebras over a polynomial algebra in two variables, in the sense of [7], a construction which includes the generalized Weyl algebras introduced by V.V. Bavula in [1]. In [8] the authors determine a basis for  $\mathcal{H}(f)$  over  $\mathbb{C}$ , compute the center of  $\mathcal{H}(f)$ , solve the isomorphism problem for this family of algebras and classify all the finite-dimensional irreducible representations of  $\mathcal{H}(f)$ .

In this note we show that  $\mathcal{H}(f)$  is (right or left) Noetherian if and only if  $\deg f = 1$  and that  $\mathcal{H}(f)$  is isomorphic to a generalized down-up algebra if and only if  $\deg f \leq 1$ . For this reason, we then concentrate on the case where  $\deg f > 1$  and determine the locally nilpotent and the locally finite derivations of  $\mathcal{H}(f)$ , all  $\mathbb{Z}$ -gradings of  $\mathcal{H}(f)$  and describe the automorphism group of  $\mathcal{H}(f)$ . In particular, we obtain the following results in case  $\deg f > 1$ :

- (i)  $\mathcal{H}(f)$  in neither right nor left Noetherian (Proposition 2.4);
- (ii)  $\mathcal{H}(f)$  admits a unique (up to an integer multiple) nontrivial  $\mathbb{Z}$ -grading, in which  $x$  has degree 1,  $y$  has degree  $-1$  and  $h$  has degree 0 (Corollary 4.10);
- (iii) the automorphism group of  $\mathcal{H}(f)$  is abelian: it is isomorphic to  $\mathbb{C}^* \times \mathbb{C}$ , where  $\mathbb{C}$  is a finite cyclic group whose order divides  $(\deg f) - 1$  (Theorem 5.5).

In Section 2 of the paper we review some properties of  $\mathcal{H}(f)$  which have been established in [8], determine when  $\mathcal{H}(f)$  is Noetherian and when it is isomorphic to a generalized down-up algebra, while in Section 3 we introduce a useful commutative subalgebra of  $\mathcal{H}(f)$ , which is a maximal commutative subalgebra if  $\deg f > 1$ . Assuming that  $\deg f > 1$ , we then investigate the locally finite and the locally nilpotent derivations of  $\mathcal{H}(f)$  and also its  $\mathbb{Z}$ -gradings in Section 4, and in the final section, Section 5, we describe the automorphism group of  $\mathcal{H}(f)$  and show that it is always an abelian group generated by the automorphisms which fix  $h$  and the automorphisms which fix  $x$ .

We make use of the commutator notation  $[a, b] = ab - ba$ . The sets of integers, nonnegative integers and positive integers are denoted by  $\mathbb{Z}$ ,  $\mathbb{Z}_{\geq 0}$  and  $\mathbb{Z}_{> 0}$ , respectively. The field of complex numbers is denoted by  $\mathbb{C}$ , and the multiplicative group of nonzero complex numbers is denoted by  $\mathbb{C}^*$ . For a polynomial  $g \in \mathbb{C}[h]$ ,  $\deg g$  will always denote the degree of  $g$  as a polynomial in  $h$ .

Throughout the paper,  $\sigma : \mathbb{C}[h] \rightarrow \mathbb{C}[h]$  is the algebra endomorphism given by  $\sigma(h) = f(h)$ . For any function  $\phi : X \rightarrow X$ , we will use the notation  $\phi^k$  to mean the  $k$ -th power of  $\phi$  with respect to composition. In particular,  $\phi^0$  denotes the identity on the set  $X$ .

## 2 The Noetherian property

Below we record a few results from [8] which will be useful in the course of this paper. As usual,  $Z(\mathcal{H}(f))$  denotes the center of  $\mathcal{H}(f)$ .

**Lemma 2.1** ([8, Lemma 1, Lemma 2, Theorem 4]). *Let  $f \in \mathbb{C}[h]$ . Then:*

- (a) The set  $\{x^i h^j y^k \mid i, j, k \in \mathbb{Z}_{\geq 0}\}$  is a basis of  $\mathcal{H}(f)$ .
- (b) The algebra  $\mathcal{H}(f)$  is a domain if and only if  $\deg f \geq 1$ .
- (c) The center of  $\mathcal{H}(f)$  contains the polynomial algebra  $\mathbb{C}[z]$ , where  $z = xy - h = yx - f(h)$ . If  $\deg f \neq 1$ , then  $Z(\mathcal{H}(f)) = \mathbb{C}[z]$ .

**Remarks 2.2.**

1. Identifying  $\mathcal{H}(f)$  with the ambiskew polynomial ring  $R(\mathbb{C}[h], \sigma, f(h) - h, 1)$  as in (1.2), it follows that  $\mathcal{H}(f)$  is conformal, as defined in [5, Section 2.3], and the corresponding Casimir element is precisely the central element  $z = xy - h$  defined above.
2. Suppose  $f \in \mathbb{C}$ . Then by considering the generators  $-x, y$  and  $h - f$ , we see that  $\mathcal{H}(f) \cong R(\mathbb{C}[h], \sigma, h, 1)$ , with  $\sigma = 0$ , and from [5, Theorem 7.10] we conclude that  $\mathcal{H}(f)$  is a prime ring. Thus by Lemma 2.1(b),  $\mathcal{H}(f)$  is a prime ring for any  $f \in \mathbb{C}[h]$ .
3. Since the center of  $\mathcal{H}(f)$  contains the polynomial algebra  $\mathbb{C}[z]$  and  $\mathcal{H}(f)$  has countable dimension over  $\mathbb{C}$ , it follows from Dixmier's version of Schur's Lemma that  $\mathcal{H}(f)$  is never primitive.

There is an order two anti-automorphism of  $\mathcal{H}(f)$ , denoted by  $\iota$ , that fixes  $h$  and interchanges  $x$  and  $y$ :

$$\iota : \mathcal{H}(f) \rightarrow \mathcal{H}(f), \quad x \mapsto y, \quad y \mapsto x, \quad h \mapsto h. \quad (2.3)$$

Hence  $\mathcal{H}(f)$  is isomorphic to its opposite algebra  $\mathcal{H}(f)^{\text{op}}$ .

**Proposition 2.4.** *The algebra  $\mathcal{H}(f)$  is right (respectively, left) Noetherian if and only if  $\deg f = 1$ .*

*Proof.* If  $\deg f = 1$  then  $\mathcal{H}(f)$  is a generalized Weyl algebra over a polynomial ring in two variables, and thus it is right and left Noetherian. So assume that  $\deg f \neq 1$ . In particular,  $f(h) - h$  has some root  $\alpha \in \mathbb{C}$ . Let  $F(h) = f(h + \alpha) - \alpha$ . Then  $\deg F = \deg f$  (here we assume the zero polynomial has degree 0) and  $F(h) \in h\mathbb{C}[h]$ . Moreover,  $F(h - \alpha) = f(h) - \alpha$  and then  $\mathcal{H}(f) \cong \mathcal{H}(F)$  by [8, Lemma 3]. So there is no loss in assuming that  $f(h) \in h\mathbb{C}[h]$ . By the isomorphism  $\mathcal{H}(f) \cong \mathcal{H}(f)^{\text{op}}$  it will be enough to show that  $\mathcal{H}(f)$  is not left Noetherian.

For each  $n \in \mathbb{Z}_{\geq 0}$  define the left ideal

$$I_n = \sum_{i=0}^n \mathcal{H}(f) h y^i.$$

Then  $I_n \subseteq I_{n+1}$  for all  $n \geq 0$  and we finish the proof by showing that these inclusions are strict. Note that by Lemma 2.1(a),

$$\mathcal{H}(f) = \bigoplus_{j,k \geq 0} x^j \mathbb{C}[h] y^k.$$

Given  $j, k \geq 0$  and  $g(h) \in \mathbb{C}[h]$ , we have  $x^j g(h) y^k h y^i = x^j g(h) \sigma^k(h) y^{k+i}$ . Assume, by way of contradiction, that  $h y^{n+1} \in I_n$ . Then there exist  $g_i(h) \in \mathbb{C}[h]$ ,  $i = 0, \dots, n$ , such that  $h y^{n+1} = \sum_{i=0}^n g_i(h) \sigma^{n+1-i}(h) y^{n+1}$ . It follows by Lemma 2.1(a) that

$$h = \sum_{i=0}^n g_i(h) \sigma^{n+1-i}(h). \quad (2.5)$$

As by hypothesis  $\sigma(h) = f(h) \in h\mathbb{C}[h]$ , one can deduce that  $\sigma^{n+1-i}(h) \in f(h)\mathbb{C}[h]$  for all  $0 \leq i \leq n$  and (2.5) then implies that  $h \in f(h)\mathbb{C}[h]$ , which is a contradiction since under our hypothesis either  $f(h) = 0$  or  $\deg f > 1$ . This proves that  $h y^{n+1} \notin I_n$  for any  $n \geq 0$  and hence  $\{I_n\}_{n \geq 0}$  is a strict ascending chain of left ideals of  $\mathcal{H}(f)$ .  $\square$

**Remark 2.6.** The case  $f \in \mathbb{C}$  of Proposition 2.4 follows also from [5, Corollary 7.3], which applies when  $\sigma$  is not injective. In terms of the endomorphism  $\sigma$ , Proposition 2.4 could be restated as: The algebra  $\mathcal{H}(f)$  is right (respectively, left) Noetherian if and only if  $\sigma$  is an automorphism.

We recall that a generalized down-up algebra  $L(g, r, s, \gamma)$ , given by the parameters  $g \in \mathbb{C}[H]$  and  $r, s, \gamma \in \mathbb{C}$ , is defined as the unital associative  $\mathbb{C}$ -algebra generated by  $d$ ,  $u$  and  $H$ , subject to the relations:

$$dH - rHd + \gamma d = 0, \quad Hu - ruH + \gamma u = 0, \quad du - sud + g(H) = 0.$$

Generalized down-up algebras were defined in [4] as generalizations of the down-up algebras introduced by Benkart and Roby in [2]. Generalized down-up algebras include all down-up algebras, the algebras similar to the enveloping algebra of  $\mathfrak{sl}_2$  defined by Smith [11], Le Bruyn's conformal  $\mathfrak{sl}_2$  enveloping algebras [6] and Rueda's algebras similar to the enveloping algebra of  $\mathfrak{sl}_2$  [10].

**Corollary 2.7.** *The algebra  $\mathcal{H}(f)$  is isomorphic to a generalized down-up algebra if and only if  $\deg f \leq 1$ .*

*Proof.* Suppose first that  $\deg f \leq 1$ , say  $f(h) = ah + b$  for  $a, b \in \mathbb{C}$ . Then it is straightforward to verify that  $\mathcal{H}(f) \cong L(H - f(H), a, 1, -b)$ , under an isomorphism that sends  $x$ ,  $y$  and  $h$  to  $u$ ,  $d$  and  $H$ , respectively. Conversely, suppose that  $\deg f > 1$ . Then by Proposition 2.4 and Lemma 2.1(b),  $\mathcal{H}(f)$  is a non-Noetherian domain. Hence  $\mathcal{H}(f)$  cannot be isomorphic to a generalized down-up algebra, as a generalized down-up algebra is a domain if and only if it is Noetherian, by Propositions 2.5 and 2.6 of [4].  $\square$

In view of this result, we will henceforth focus most of our attention on the generalized Heisenberg algebras  $\mathcal{H}(f)$  with  $f \in \mathbb{C}[h]$  such that  $\deg f > 1$ .

### 3 The commutative algebra $\mathcal{H}(f)_0$

In this short section we record a few useful formulas for computing in  $\mathcal{H}(f)$  and then explore an interesting commutative subalgebra of  $\mathcal{H}(f)$ .

**Lemma 3.1.** *Let  $k \in \mathbb{Z}_{\geq 0}$  and  $g \in \mathbb{C}[h]$ . Then the following hold:*

- (a)  $[y, x^k] = x^{k-1}(\sigma^k(h) - h)$ ;
- (b)  $[y^k, x] = (\sigma^k(h) - h)y^{k-1}$ ;
- (c)  $(x^k g y^k)x = x(x^k \sigma(g)y^k + x^{k-1}(\sigma^k(h) - h)g y^{k-1})$ ;
- (d)  $y(x^k g y^k) = (x^k \sigma(g)y^k + x^{k-1}(\sigma^k(h) - h)g y^{k-1})y$ ;
- (e)  $x^k g y^k$  commutes with  $x^j \tilde{g} y^j$  for all  $\tilde{g} \in \mathbb{C}[h]$  and all  $j \in \mathbb{Z}_{\geq 0}$ .

*Proof.* Parts (a) and (b) have been established in [5], formulas (6a)–(6b). We prove part (c) using (b):

$$\begin{aligned} (x^k g y^k)x &= x^k g x y^k + x^k g [y^k, x] \\ &= x^{k+1} \sigma(g) y^k + x^k g (\sigma^k(h) - h) y^{k-1} \\ &= x(x^k \sigma(g) y^k + x^{k-1}(\sigma^k(h) - h)g y^{k-1}). \end{aligned}$$

Formula (d) follows from applying the anti-automorphism  $\iota$  of (2.3) to (c).

Finally, we prove (e) by induction on  $k$ , the case  $k = 0$  being trivial:

$$g(x^j \tilde{g} y^j) = x^j \sigma^j(g) \tilde{g} y^j = x^j \tilde{g} \sigma^j(g) y^j = (x^j \tilde{g} y^j)g.$$

Now suppose (e) holds for a certain  $k \geq 0$ . Thus we have:

$$\begin{aligned} (x^{k+1} g y^{k+1})(x^j \tilde{g} y^j) &= (x^{k+1} g y^k) y (x^j \tilde{g} y^j) \\ &= x(x^k g y^k)(x^j \sigma(\tilde{g}) y^j) + x^{j-1}(\sigma^j(h) - h) \tilde{g} y^{j-1} y \quad \text{by (d)} \\ &= x(x^j \sigma(g) y^k + x^{j-1}(\sigma^j(h) - h) \tilde{g} y^{j-1})(x^k g y^k) y \quad (*) \\ &= (x^j \tilde{g} y^j) x (x^k g y^k) y \quad \text{by (c)} \\ &= (x^j \tilde{g} y^j)(x^{k+1} g y^{k+1}), \end{aligned}$$

where (\*) follows from the induction hypothesis. So (e) holds for all  $k \in \mathbb{Z}_{\geq 0}$ .  $\square$

There is an obvious grading of  $\mathcal{H}(f)$  relative to which  $x$  has degree 1,  $y$  has degree  $-1$  and  $h$  has degree 0. We denote the corresponding homogeneous subspaces by  $\mathcal{H}(f)_\ell$ , for  $\ell \in \mathbb{Z}$ , so that

$$\mathcal{H}(f) = \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}(f)_\ell, \quad \text{with} \quad \mathcal{H}(f)_\ell = \bigoplus_{i-k=\ell} \mathbb{C} x^i \mathbb{C}[h] y^k. \quad (3.2)$$

We call this the *standard grading* of  $\mathcal{H}(f)$ , and, whenever we mention a homogeneous component or element of  $\mathcal{H}(f)$ , we will always be referring to this standard grading.

The subalgebra  $\mathcal{H}(f)_0$  has basis  $\{x^k h^j y^k \mid k, j \geq 0\}$  and  $\mathcal{H}(f)_\ell = x^\ell \mathcal{H}(f)_0$  if  $\ell \geq 0$ ;  $\mathcal{H}(f)_\ell = \mathcal{H}(f)_0 y^{-\ell}$  if  $\ell \leq 0$ . Thus we have the decomposition

$$\mathcal{H}(f) = \bigoplus_{\ell > 0} x^\ell \mathcal{H}(f)_0 \oplus \mathcal{H}(f)_0 \oplus \bigoplus_{\ell > 0} \mathcal{H}(f)_0 y^\ell.$$

**Proposition 3.3.** *The subalgebra  $\mathcal{H}(f)_0$  is commutative. If  $\deg f > 1$ , then  $\mathcal{H}(f)_0$  is a maximal commutative subalgebra of  $\mathcal{H}(f)$  which strictly contains  $\mathbb{C}[z, h]$ , the polynomial subalgebra of  $\mathcal{H}(f)$  generated by  $h$  and the central element  $z = xy - h$ .*

*Proof.* The first statement is a direct consequence of Lemma 3.1(e). Assume now that  $\deg f > 1$ . Then  $\sigma$  is injective and has infinite order. For any  $i, k \in \mathbb{Z}_{\geq 0}$  and  $g \in \mathbb{C}[h]$ ,  $[h, x^i g y^k] = x^i (\sigma^i(h) - \sigma^k(h)) g y^k$ . Hence, if  $g \neq 0$ , we deduce that  $[h, x^i g y^k] = 0 \iff i = k$ , and from this it is straightforward to conclude that  $\mathcal{H}(f)_0$  is the centralizer of  $h$ , hence a maximal commutative subalgebra of  $\mathcal{H}(f)$ .

The commuting elements  $h$  and  $z$  are homogeneous of degree 0 and are easily seen to be algebraically independent, as  $z^k - x^k y^k$  is in the span of  $\{x^i g y^i \mid i < k, g \in \mathbb{C}[h]\}$ . Suppose, by contradiction, that there exist  $g_k \in \mathbb{C}[h]$  such that  $xhy = \sum_{k \geq 0} g_k z^k$ . Then by the argument above we must have  $g_k = 0$  for all  $k > 1$  and  $\sigma(g_1) = h$ , which is possible only if  $\deg f = 1$ . Therefore  $xhy \in \mathcal{H}(f)_0 \setminus \mathbb{C}[z, h]$ .  $\square$

By Lemma 3.1(c)–(d), it is possible to extend  $\sigma$  to a  $\mathbb{C}$ -linear endomorphism  $\tilde{\sigma}$  of  $\mathcal{H}(f)_0$  so that  $\tilde{\sigma}(x^k g y^k) = x^k \sigma(g) y^k + x^{k-1} (\sigma^k(h) - h) g y^{k-1}$ , for all  $k \in \mathbb{Z}_{\geq 0}$  and  $g \in \mathbb{C}[h]$ . For simplicity, we still denote this endomorphism by  $\sigma$  instead of  $\tilde{\sigma}$ . By Lemma 3.1(c)–(d) and Lemma 2.1(a),  $\sigma$  is defined by the relations:

$$\theta x = x \sigma(\theta), \quad y \theta = \sigma(\theta) y, \quad \text{for all } \theta \in \mathcal{H}(f)_0. \quad (3.4)$$

In particular, (3.4) implies that  $\sigma$  is an algebra endomorphism of  $\mathcal{H}(f)_0$ .

## 4 locally finite derivations of $\mathcal{H}(f)$ when $\deg f > 1$

Henceforth we will assume that  $\deg f > 1$ . By Corollary 2.7 we are assuming that  $\mathcal{H}(f)$  is not a generalized down-up algebra. Most of our subsequent results do not hold if  $\deg f \leq 1$ .

Our goal in this section is to determine all locally finite derivations of  $\mathcal{H}(f)$ . In particular, we will classify all  $\mathbb{Z}$ -gradings of  $\mathcal{H}(f)$  and show that  $\mathcal{H}(f)$  has no nontrivial locally nilpotent derivations. Our methods are akin to those used in [12].

Let  $\delta$  be a  $\mathbb{C}$ -linear endomorphism of  $\mathcal{H}(f)$ . We recall the following standard definitions:

- $\delta$  is a derivation of  $\mathcal{H}(f)$  if  $\delta(ab) = \delta(a)b + a\delta(b)$ ;
- $\delta$  is locally finite if for every  $a \in \mathcal{H}(f)$  the  $\mathbb{C}$ -linear span of  $\{\delta^k(a) \mid k \in \mathbb{Z}_{\geq 0}\}$  is finite dimensional;
- $\delta$  is locally nilpotent if for every  $a \in \mathcal{H}(f)$  there is  $k \in \mathbb{Z}_{\geq 0}$  such that  $\delta^k(a) = 0$ ;

- $\delta$  is homogeneous of degree  $r \in \mathbb{Z}$  if  $\delta(\mathcal{H}(f)_\ell) \subseteq \mathcal{H}(f)_{\ell+r}$  for all  $\ell \in \mathbb{Z}$ .

Assume  $\delta$  is any derivation of  $\mathcal{H}(f)$ . Since  $\mathcal{H}(f)$  is finitely generated, there exist homogeneous derivations  $\delta_1, \dots, \delta_k$  of strictly increasing degrees such that  $\delta = \delta_1 + \dots + \delta_k$ . Moreover, as seen in [12, Lemma 1.1], if  $\delta$  is locally finite, then so are  $\delta_1$  and  $\delta_k$ , and if  $\delta_1$  (respectively,  $\delta_k$ ) is of nonzero degree, then it must be locally nilpotent.

We need one final definition. Given a locally nilpotent derivation  $\delta$  and  $a \in \mathcal{H}(f)$ , define

$$\deg_\delta(a) = \max\{k \in \mathbb{Z}_{\geq 0} \mid \delta^k(a) \neq 0\} \quad \text{if } a \neq 0;$$

define also  $\deg_\delta(0) = -\infty$ . It can be easily checked (see for example [9]) that for  $a, b \in \mathcal{H}(f)$ ,  $\deg_\delta(a + b) \leq \max\{\deg_\delta(a), \deg_\delta(b)\}$ , with equality if  $\deg_\delta(a) \neq \deg_\delta(b)$ . Since  $\mathcal{H}(f)$  is a domain and  $\mathbb{C}$  has characteristic 0, we also have, from the Leibniz rule,  $\deg_\delta(ab) = \deg_\delta(a) + \deg_\delta(b)$ . In particular,  $\ker \delta$  is factorially closed: if  $\delta(ab) = 0$  for some nonzero  $a, b \in \mathcal{H}(f)$ , then  $\delta(a) = 0 = \delta(b)$ .

**Lemma 4.1.** *Assume that  $\deg f > 1$ . Then all locally finite derivations of  $\mathcal{H}(f)$  are homogeneous of degree 0.*

*Proof.* Let  $\delta$  be a locally finite derivation of  $\mathcal{H}(f)$ . By the above decomposition  $\delta = \delta_1 + \dots + \delta_k$  of  $\delta$  into homogeneous derivations of strictly increasing degrees, it will be enough to show that there are no nonzero homogeneous locally nilpotent derivations of  $\mathcal{H}(f)$  of degree  $r \neq 0$ .

So assume  $\delta$  is a homogeneous locally nilpotent derivation of  $\mathcal{H}(f)$  of degree  $r \neq 0$ . Let  $d = \deg_\delta(h)$  and suppose that  $d > 0$ . Then  $\deg_\delta(f(h)) = d \deg f$  and the relation  $hx = xf(h)$  yields

$$d + \deg_\delta(x) = \deg_\delta(x) + d \deg f,$$

so  $\deg f = 1$ , which contradicts our hypothesis. Hence  $d = 0$  and  $\delta(h) = 0$ .

By replacing  $\delta$  with  $\iota \delta \iota^{-1}$ , where  $\iota$  is the anti-automorphism defined in (2.3), we can assume that  $r > 0$ . Then  $\ker \delta$  contains some nonzero homogeneous element of positive degree. Since elements of  $\mathcal{H}(f)$  of positive degree lie in  $x\mathcal{H}(f)$  and  $\ker \delta$  is factorially closed, we deduce that  $\delta(x) = 0$ .

Any derivation maps the center of an algebra into itself, so  $\delta$  restricts to a locally nilpotent derivation of  $\mathbb{C}[z]$ , by Lemma 2.1(c), and thus  $\delta(z) \in \mathbb{C}$ . On the other hand, since  $z = xy - h$  is homogeneous of degree 0 and  $\delta$  has positive degree, it must be that  $\delta(z) = 0$ , and from  $0 = \delta(z) = x\delta(y)$ , we conclude that  $\delta(y) = 0$ . Then  $\delta = 0$  and the lemma is proved.  $\square$

The next theorem, our main result on derivations of  $\mathcal{H}(f)$  when  $\deg f > 1$ , shows that the space of locally finite derivations of  $\mathcal{H}(f)$  is one-dimensional over  $\mathbb{C}$ , spanned by the derivation  $\partial$  defined by

$$\partial(x^i h^j y^k) = (i - k)x^i h^j y^k, \quad \text{for all } i, j, k \in \mathbb{Z}_{\geq 0}. \quad (4.2)$$

**Theorem 4.3.** *Assume that  $\deg f > 1$ . If  $\delta$  is a locally finite derivation of  $\mathcal{H}(f)$ , then there is  $\lambda \in \mathbb{C}$  such that  $\delta(x) = \lambda x$ ,  $\delta(y) = -\lambda y$  and  $\delta(h) = 0$ .*

*Proof.* Let  $\delta$  be a locally finite derivation of  $\mathcal{H}(f)$ . By Lemma 4.1, we know that  $\delta$  is homogeneous of degree 0, so there are  $\theta_x, \theta_h, \theta_y \in \mathcal{H}(f)_0$  so that

$$\delta(x) = x\theta_x, \quad \delta(h) = \theta_h, \quad \text{and} \quad \delta(y) = \theta_y y.$$

In particular, since  $h$  commutes with  $\theta_h$ , we have  $\delta(g(h)) = g'(h)\theta_h$  for all  $g(h) \in \mathbb{C}[h]$ , where  $g'(h)$  denotes the derivative of  $g(h)$  with respect to  $h$ .

*Claim 1:*  $\theta_h = 0$  and  $\theta_x + \theta_y = 0$ .

*Proof of Claim 1:* Write

$$\theta_h = \sum_{k \geq 0} x^k g_k(h) y^k, \tag{4.4}$$

with  $g_k(h) \in \mathbb{C}[h]$  and  $g_k(h) = 0$  except for finitely many indices  $k$ .

As observed in the proof of Lemma 4.1,  $\delta$  restricts to a locally finite derivation of  $\mathbb{C}[z]$ , the center of  $\mathcal{H}(f)$ , and thus  $\delta(z) \in \mathbb{C} \oplus \mathbb{C}z$ , say  $\delta(z) = \mu z - \lambda$ , with  $\lambda, \mu \in \mathbb{C}$ . Since  $\mu z - \lambda = \delta(xy - h) = x(\theta_x + \theta_y)y - \theta_h$ , we have

$$\theta_h = x(\theta_x + \theta_y)y - \mu z + \lambda = x(\theta_x + \theta_y - \mu)y + \mu h + \lambda. \tag{4.5}$$

In particular,  $g_0(h) = \mu h + \lambda$ .

We now apply  $\delta$  to the relation  $yh = f(h)y$  and get  $\theta_y y h + y \theta_h = f'(h)\theta_h y + f(h)\theta_y y$ . As  $h$  and  $\theta_y$  commute, and  $y \theta_h = \sigma(\theta_h)y$ , by (3.4), we obtain

$$\sigma(\theta_h) = f'(h)\theta_h. \tag{4.6}$$

Now combining (4.4) and (4.6) we deduce that, for every  $k \geq 0$ :

$$\sigma^k(f'(h))g_k(h) = \sigma(g_k(h)) + (\sigma^{k+1}(h) - h)g_{k+1}(h). \tag{4.7}$$

Setting  $k = 0$  in (4.7) we obtain  $(f(h) - h)g_1(h) = f'(h)g_0(h) - \sigma(g_0(h))$ . Since we have already established that  $\deg g_0 \leq 1$ , we deduce now from the latter equation that  $\deg(f(h) - h)g_1(h) \leq \deg f$ , and thus  $g_1 \in \mathbb{C}$ . Combining this with the  $k = 1$  case of (4.7),  $\sigma(f'(h))g_1(h) = \sigma(g_1(h)) + (\sigma(f(h)) - h)g_2(h)$ , yields  $g_2 = 0$ , and in turn the latter gives  $g_k = 0$  for all  $k \geq 2$ . Using again the relation  $\sigma(f'(h))g_1(h) = \sigma(g_1(h)) + (\sigma(f(h)) - h)g_2(h)$  with  $g_2 = 0$  and  $g_1 \in \mathbb{C}$  gives  $g_1 = 0$ . Therefore we have

$$\sigma(g_0) = f'(h)g_0. \tag{4.8}$$

Suppose  $g_0 \neq 0$ , and let  $a$  be the leading coefficient of  $f(h)$ . Then  $\mu \neq 0$  and comparing leading coefficients in (4.8) yields  $\mu a = a(\deg f)\mu$ , whence  $\deg f = 1$ , which is a contradiction. Thus  $g_0 = 0$ .

From the above we conclude that  $\theta_h = \sum_{k \geq 0} x^k g_k(h) y^k = 0$  and finally by (4.5) we get  $\theta_x + \theta_y = 0$ , establishing Claim 1.

So far we have shown that

$$\delta(x) = x\theta_x, \quad \delta(h) = 0, \quad \text{and} \quad \delta(y) = -\theta_x y,$$



so it remains to be inferred that  $\theta_x \in \mathbb{C}$ .

*Claim 2:*  $\delta(\theta) = 0$ , for all  $\theta \in \mathcal{H}(f)_0$ .

*Proof of Claim 2:* Since  $\delta(g) = 0$  for all  $g \in \mathbb{C}[h]$ , it suffices to show that if  $\theta \in \mathcal{H}(f)_0$  and  $\delta(\theta) = 0$ , then also  $\delta(x\theta y) = 0$ . This follows easily using the fact that  $\mathcal{H}(f)_0$  is commutative, as proved in Proposition 3.3.

From Claim 2 it follows that, for all  $k \geq 0$ ,  $\delta(\theta_x^k) = 0$ , which implies that  $\delta^k(x) = x\theta_x^k$ . As  $\delta$  is locally finite, the span of  $\{\theta_x^k \mid k \in \mathbb{Z}_{\geq 0}\}$  must then be finite dimensional. This is possible only if  $\theta_x \in \mathbb{C}$ , thus finishing the proof of the theorem.  $\square$

Since locally nilotent derivations are locally finite, we derive the following corollary.

**Corollary 4.9.** *Assume that  $\deg f > 1$ . Then  $\mathcal{H}(f)$  has no nonzero locally nilpotent derivations.*

Suppose that  $\mathcal{H}(f) = \bigoplus_{\alpha \in \mathbb{C}} V_\alpha$  is a grading. Define the  $\mathbb{C}$ -linear endomorphism  $\delta$  of  $\mathcal{H}(f)$  by  $\delta(v_\alpha) = \alpha v_\alpha$  for all  $v_\alpha \in V_\alpha$  and all  $\alpha \in \mathbb{C}$ . It is immediate to check that  $\delta$  is a diagonalizable derivation of  $\mathcal{H}(f)$  whose eigenvalues are those  $\alpha \in \mathbb{C}$  such that  $V_\alpha \neq (0)$ . Conversely, if  $\delta$  is a diagonalizable derivation, then  $\delta$  determines a grading where  $V_\alpha$  is the  $\alpha$ -eigenspace of  $\delta$ . Furthermore, diagonalizable derivations are clearly locally finite.

Thus, we deduce from Theorem 4.3 that, except for the trivial grading in which every element of  $\mathcal{H}(f)$  has degree 0,  $\mathcal{H}(f)$  only admits the standard grading defined in (3.2), up to scaling by some integer. More precisely, we have:

**Corollary 4.10.** *Assume that  $\deg f > 1$ . Then for any  $\mathbb{Z}$ -grading of  $\mathcal{H}(f)$ , there is an integer  $\ell \in \mathbb{Z}$  such that, relative to that grading,  $x$  has degree  $\ell$ ,  $y$  has degree  $-\ell$  and  $h$  has degree 0.*

## 5 Automorphisms of $\mathcal{H}(f)$ when $\deg f > 1$

When  $\deg f = 1$  the algebra  $\mathcal{H}(f)$  is a Noetherian generalized down-up algebra, by Corollary 2.7, and the automorphisms of the latter have been investigated in [3]. We continue to assume that  $\deg f > 1$  and note again that our results do not generalize to the cases with  $\deg f \leq 1$ .

Since  $\mathcal{H}(f)$  has no nonzero locally nilpotent derivations, it seems natural to conjecture that the automorphism group of  $\mathcal{H}(f)$  is somewhat small. However, over  $\mathbb{C}$  we can consider also the exponential of a diagonalizable derivation. Specifically, let  $c \in \mathbb{C}$  and let  $\partial$  be the derivation of  $\mathcal{H}(f)$  defined in (4.2). Then the expression

$$\exp(c\partial) := \sum_{k=0}^{\infty} \frac{(c\partial)^k}{k!}$$

defines an automorphism of  $\mathcal{H}(f)$  satisfying

$$\exp(c\partial)(x) = \sum_{k=0}^{\infty} \frac{c^k}{k!} x = \exp(c)x, \quad \exp(c\partial)(y) = \exp(-c)y, \quad \exp(c\partial)(h) = h,$$

with inverse  $\exp(-c\partial)$ .

The above motivates the following definition. For each  $\lambda \in \mathbb{C}^*$ , let  $\phi_\lambda$  be the automorphism of  $\mathcal{H}(f)$  defined by

$$\phi_\lambda(x) = \lambda x, \quad \phi_\lambda(y) = \lambda^{-1}y, \quad \phi_\lambda(h) = h. \quad (5.1)$$

The group of algebra automorphisms of  $\mathcal{H}(f)$  will be denoted by  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ . We have a first description of  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  below.

**Proposition 5.2.** *Assume  $\deg f > 1$ . Then the following hold:*

- (a) *Any automorphism of  $\mathcal{H}(f)$  restricts to an automorphisms of  $\mathbb{C}[h]$ , and  $x$  and  $y$  are eigenvectors.*
- (b)  *$\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(h) = h\} = \{\phi_\lambda \mid \lambda \in \mathbb{C}^*\} \cong \mathbb{C}^*$ , and this is a central subgroup of  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ .*
- (c)  *$\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is a finite cyclic subgroup whose order divides  $(\deg f) - 1$ .*

*Proof.* Let  $\phi$  be an automorphism of  $\mathcal{H}(f)$ . Then as argued in Claim 4 of the proof of [8, Theorem 5], the relation  $\phi(h)\phi(x) = \phi(x)f(\phi(h))$  with  $\deg f > 1$  implies that  $\phi(h) \in \mathbb{C}[h]$ ; applying this result to  $\phi^{-1}$  gives that  $\phi(h) = ah + b$ , for some  $a, b \in \mathbb{C}$  with  $a \neq 0$ .

Now writing  $\phi(x)$  as a sum of terms of the form  $x^i g_{i,j} y^j$  with  $i, j \in \mathbb{Z}_{\geq 0}$  and  $g_{i,j} \in \mathbb{C}[h]$ , and comparing the corresponding expressions for  $\phi(h)\phi(x)$  and  $\phi(x)f(\phi(h))$ , we obtain  $\phi(x) \in \mathcal{H}(f)_1$ . Similarly,  $\phi(y) \in \mathcal{H}(f)_{-1}$ , so  $\phi$  is homogeneous of degree 0. Thus, there exist  $\theta_x, \theta_y \in \mathcal{H}(f)_0$  such that  $\phi(x) = x\theta_x$  and  $\phi(y) = \theta_y y$ . Applying the same reasoning to  $\phi^{-1}$ , we deduce that  $\theta_x, \theta_y \in \mathbb{C}^*$ , which proves (a).

Now assume  $\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  and  $\phi(h) = h$ . By (a) there exist  $\lambda, \mu \in \mathbb{C}^*$  such that  $\phi(x) = \lambda x$  and  $\phi(y) = \mu y$ . Applying  $\phi$  to the relation  $[y, x] = f(h) - h$  yields  $\lambda\mu = 1$ , so  $\phi = \phi_\lambda$ . This proves the equality in (b), and the isomorphism  $\{\phi_\lambda \mid \lambda \in \mathbb{C}^*\} \cong \mathbb{C}^*$  is clear, as  $\phi_\lambda \circ \phi_\mu = \phi_{\lambda\mu}$  for all  $\lambda, \mu \in \mathbb{C}^*$ .

Next, we show that the subgroup  $\{\phi_\lambda \mid \lambda \in \mathbb{C}^*\}$  is central in  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ . Let  $\lambda \in \mathbb{C}^*$ , and suppose  $\psi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  is arbitrary. By (a) we know that  $\psi(h) \in \mathbb{C}[h]$ , which implies that  $\phi_\lambda \circ \psi(h) = \psi \circ \phi_\lambda(h)$ . But as  $x$  and  $y$  are eigenvalues for any automorphism of  $\mathcal{H}(f)$ ,  $\phi_\lambda \circ \psi$  and  $\psi \circ \phi_\lambda$  also agree on these generators, and thus  $\phi_\lambda \circ \psi = \psi \circ \phi_\lambda$ .

To prove part (c), suppose that  $\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  and  $\phi(x) = x$ . We know already that  $\phi(h) = ah + b$  and  $\phi(y) = cy$ , for some  $a, b, c \in \mathbb{C}$  with  $a, c \neq 0$ . Then  $xf(ah + b) = x\phi(f(h)) = \phi(h)x = (ah + b)x = x(af(h) + b)$ , and we obtain

$$f(ah + b) = af(h) + b. \quad (5.3)$$

Therefore,

$$\begin{aligned} c(f(h) - h) &= c[y, x] = \phi([y, x]) = \phi(f(h) - h) = af(h) + b - (ah + b) \\ &= a(f(h) - h), \end{aligned}$$

and we conclude that  $c = a$ .

Write  $f(h) = \sum_{k=0}^n a_k h^k$ , where  $n = \deg f$  and  $a_k \in \mathbb{C}$ . Applying the derivation  $\frac{d}{dh}$  to (5.3)  $n - 1$  times yields  $a^{n-1} f^{(n-1)}(ah + b) = a f^{(n-1)}(h)$ , as  $n - 1 \geq 1$ . As  $f^{(n-1)}(h) = (n - 1)!(na_n h + a_{n-1})$ , we obtain

$$a^{n-1} = 1 \quad \text{and} \quad b = \frac{(a - 1)a_{n-1}}{na_n}. \quad (5.4)$$

Let  $U_{n-1} = \{\zeta \in \mathbb{C}^* \mid \zeta^{n-1} = 1\}$  be the cyclic group of order  $n - 1$ , and define a map

$$\Psi : \{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\} \longrightarrow U_{n-1}, \quad \phi \mapsto a, \quad \text{where } \phi(h) = ah + b.$$

Then  $\Psi$  is well defined by (5.4), and it is a group homomorphism. If  $\Psi(\phi) = 1$  for some  $\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  with  $\phi(x) = x$ , then the above shows that  $\phi(y) = y$  and  $\phi(h) = h + b$ . Again by (5.4) we deduce that  $b = 0$ , so  $\phi$  is the identity on  $\mathcal{H}(f)$ . This shows that  $\Psi$  is an injective group homomorphism and thus  $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is isomorphic to a subgroup of  $U_{n-1}$ ; hence it is a finite cyclic group whose order divides  $n - 1$ .  $\square$

It is now an easy matter to determine the structure of  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ . The symbol  $\dot{\times}$  used below denotes the internal direct product of subgroups of a group.

**Theorem 5.5.** *Assume  $\deg f > 1$ . Then*

$$\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) = \{\phi_{\lambda} \mid \lambda \in \mathbb{C}^*\} \dot{\times} \{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\} \quad (5.6)$$

*is an abelian group, where:*

- $\{\phi_{\lambda} \mid \lambda \in \mathbb{C}^*\} \cong \mathbb{C}^*$  and  $\phi_{\lambda}$  is defined in (5.1);
- $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is a finite cyclic group whose order divides  $(\deg f) - 1$  and which, as a set, can be identified with  $\{(a, b) \in \mathbb{C}^* \times \mathbb{C} \mid f(ah + b) = af(h) + b\}$  via the correspondence  $\phi \mapsto (a, b)$ , where  $\phi(h) = ah + b$ .

*Proof.* Since we have already seen in Proposition 5.2 that  $\{\phi_{\lambda} \mid \lambda \in \mathbb{C}^*\}$  is central, in order to prove the direct product decomposition in (5.6), it remains to show that the two subgroups have trivial intersection, which is clear, and generate  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ . Let  $\psi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$ . Then there is  $\lambda \in \mathbb{C}^*$  such that  $\psi(x) = \lambda x$ , whence  $\phi_{\lambda}^{-1} \circ \psi \in \{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$ , and this shows the latter claim. Moreover, since  $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is abelian, by Proposition 5.2(c), the group  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f))$  must also be abelian.

The remaining parts of the theorem have already been proved, except for the observation that  $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  can be identified with the set  $\{(a, b) \in \mathbb{C}^* \times \mathbb{C} \mid f(ah + b) = af(h) + b\}$ . Indeed, if  $\phi(x) = x$ , then we have seen in the proof of Proposition 5.2 that  $\phi(h) = ah + b$  and  $\phi(y) = ay$ , for some  $a, b \in \mathbb{C}$  with  $a \neq 0$ , and (5.3) must hold. This shows that the correspondence  $\phi \mapsto (a, b)$  is well defined and one-to-one. Conversely, given  $(a, b) \in \mathbb{C}^* \times \mathbb{C}$  satisfying  $f(ah + b) = af(h) + b$ , it is routine to check that there is an automorphism of  $\mathcal{H}(f)$  defined by the conditions  $\phi(x) = x$ ,  $\phi(y) = ay$ ,  $\phi(h) = ah + b$ , and this shows the correspondence is onto.  $\square$

**Remark 5.7.** Any pair  $(a, b) \in \mathbb{C}^* \times \mathbb{C}$  satisfying  $f(ah + b) = af(h) + b$  must also satisfy (5.4), where  $n = \deg f$ , although the conditions in (5.4) are not sufficient (see the examples below). Thus, for each  $(n - 1)$ -th root of unity  $a$ , the corresponding scalar  $b$  is determined by (5.4), but one still needs to check the relation  $f(ah + b) = af(h) + b$  for the pair  $(a, b)$ .

**Examples 5.8.**

- (a) If  $\deg f = 2$ , then  $n = 2$  in (5.4), so  $a = 1$  and  $b = 0$ , and the pair  $(1, 0)$  corresponds to the identity map. It follows that the group  $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is trivial and  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^*$ .
- (b) Let  $f(h) = h^3 + h$ . Then  $n = 3$  in (5.4), so  $a = \pm 1$ . If  $a = 1$ , then  $b = 0$ , and the corresponding automorphism is the identity. If  $a = -1$ , then  $b = 0$ , and in fact  $f(-h) = -f(h)$ . Therefore there is an automorphism  $\phi$  of  $\mathcal{H}(f)$  such that  $\phi(x) = x$ ,  $\phi(y) = -y$ ,  $\phi(h) = -h$ , and  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^* \times \mathbb{Z}_2$ .
- (c) Let  $f(h) = h^3 + h + 1$ . Then  $n = 3$  in (5.4), so  $a = \pm 1$ . If  $a = -1$ , then (5.4) yields  $b = 0$ , but  $f(-h) \neq -f(h)$ , so the group  $\{\phi \in \text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \mid \phi(x) = x\}$  is trivial and  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^*$ .
- (d) Let  $f(h) = h^n$ , for  $n > 1$ . Then (5.4) says that  $a$  is a  $(n - 1)$ -th root of unity and  $b = 0$ . Moreover,  $f(ah) = a^n f(h) = af(h)$  for any  $(n - 1)$ -th root of unity  $a$ . Hence  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^* \times \mathbb{Z}_{n-1}$ .
- (e) Let  $f(h) = h^n + 1$ , for  $n > 1$ . Then, as before,  $a$  is a  $(n - 1)$ -th root of unity and  $b = 0$ . However, in this case,  $f(ah) = ah^n + 1$  whereas  $af(h) = ah^n + a$ , so equality holds if and only if  $a = 1$ . Hence  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^*$ .
- (f) Let  $f(h) = h^n + h^{k+1}$ , for some  $n \geq 4$ , and take any  $1 \leq k < n - 1$  such that  $k \mid n - 1$ . Then  $a$  is a  $(n - 1)$ -th root of unity and  $b = 0$ . In this case  $f(ah) = ah^n + a^{k+1}h^{k+1}$  whereas  $af(h) = ah^n + ah^{k+1}$ , so equality holds if and only if  $a^k = 1$ . By the hypothesis that  $k \mid n - 1$ , we deduce that  $\text{Aut}_{\mathbb{C}}(\mathcal{H}(f)) \cong \mathbb{C}^* \times \mathbb{Z}_k$ .

**References**

- [1] V. V. Bavula. Generalized Weyl algebras and their representations. *Algebra i Analiz*, 4(1):75–97, 1992.
- [2] G. Benkart and T. Roby. Down-up algebras. *J. Algebra*, 209(1):305–344, 1998.
- [3] Paula A. A. B. Carvalho and Samuel A. Lopes. Automorphisms of generalized down-up algebras. *Comm. Algebra*, 37(5):1622–1646, 2009.
- [4] Thomas Cassidy and Brad Shelton. Basic properties of generalized down-up algebras. *J. Algebra*, 279(1):402–421, 2004.

- [5] David A. Jordan. Down-up algebras and ambiskew polynomial rings. *J. Algebra*, 228(1):311–346, 2000.
- [6] Lieven Le Bruyn. Conformal  $\mathfrak{sl}_2$  enveloping algebras. *Comm. Algebra*, 23(4):1325–1362, 1995.
- [7] Rencai Lü, Volodymyr Mazorchuk, and Kaiming Zhao. Simple weight modules over weak generalized Weyl algebras. *J. Pure Appl. Algebra*, 219(8):3427–3444, 2015.
- [8] Rencai Lü and Kaiming Zhao. Finite-dimensional simple modules over generalized Heisenberg algebras. *Linear Algebra Appl.*, 475:276–291, 2015.
- [9] L. Makar-Limanov. Locally nilpotent derivations, a new ring invariant and applications. Available at <http://math.wayne.edu/~lml/lmlnotes.pdf>.
- [10] Sonia Rueda. Some algebras similar to the enveloping algebra of  $\mathfrak{sl}(2)$ . *Comm. Algebra*, 30(3):1127–1152, 2002.
- [11] S. P. Smith. A class of algebras similar to the enveloping algebra of  $\mathfrak{sl}(2)$ . *Trans. Amer. Math. Soc.*, 322(1):285–314, 1990.
- [12] Mariano Suárez-Alvarez and Quimey Vivas. Automorphisms and isomorphisms of quantum generalized Weyl algebras. *J. Algebra*, 424:540–552, 2015.

SAMUEL A. LOPES

*CMUP, Faculdade de Ciências, Universidade do Porto*

*Rua do Campo Alegre 687*

*4169-007 Porto, Portugal*

`slopes@fc.up.pt`