







Commentary: A Position Statement on Sexual Orientation Conversion Therapies by Members of the Board of Directors of the Portuguese Society of Clinical Sexology (SPSC)

Patricia M. Pascoal^{a,b,c,d} , Gabriela Moita^b, Tiago R. Marques^{b,e,f}, Ana Carvalheira^{b,g} , Sandra Vilarinho^{b,c}, Jorge Cardoso^{b,h} , Rui F. Carvalho^b, M. Joana Almeida^{b,i} and Pedro J. Nobre^c 

^aCICPSI, Faculdade de Psicologia, Universidade de Lisboa, Lisboa, Portugal; ^bSociedade Portuguesa de Sexologia Clínica, Lisboa, Portugal; ^cCPUP, Faculdade de Psicologia e Ciências da Educação da Universidade do Porto, Porto, Portugal; ^dEscola de Psicologia e Ciências da Vida, Universidade Lusófona de Humanidades e Tecnologias, Lisboa, Portugal; ^eImperial College London Faculty of Medicine, Psychiatric Imaging Group, London, United Kingdom of Great Britain and Northern Ireland; ^fKing's College London, Department of Psychosis, Institute of Psychiatry, Psychology and Neuroscience, London, United Kingdom of Great Britain and Northern Ireland; ^gWilliam James Research Center, Department of Clinical Psychology, ISPA Instituto Universitário, Lisbon, Portugal; ^hEgas Moniz Interdisciplinary Research Centre, Caparica, Portugal; ⁱCentro de Desenvolvimento-Diferenças, Lisboa, Portugal

Studies carried out in the most diverse scientific fields (in particular, medicine and psychology) refute any association between homosexuality and pathology. In the social sciences, the evidence points to the serious harm of such assumptions, both personally and socially.

In 1973, the American Psychiatric Association (APA), recognizing this evidence, removed homosexuality from its list of mental illnesses and went on to explicitly condemn any attempt of what some call *reorientation* or *conversion therapy* (Drescher, 2015).

Two decades later, in 1992, the World Health Organization took the same position (Cochran et al., 2014).

Nonheterosexual sexual orientation is not a disease, disorder, or clinical syndrome. It does not, therefore, make sense for mental health technicians, or others, to treat sexual orientation using techniques and procedures that aim to improve people's lives and should not serve personal moral convictions (Greene, 2007). Moreover, improper use of these procedures may aggravate the suffering of those seeking help on grounds of sexual orientation (APA, 2000; Pan American Health Organization & World Health Organization, 2012).

It is true that non-heterosexual sexual orientation is often associated with psychological distress caused by social and family exclusion,

bullying, social homophobia, violence, professional discrimination, heterosexism and internalized homophobia (e.g., Hall, 2018; Ross et al., 2018).

Moreover, the effects of homophobia are felt at different moments in the life cycle, and especially in the periods of psychological, emotional, and social transition, and thus of greater vulnerability. For this reason, promoting a nonpathologizing approach and reducing personal suffering, if any, of those who are homo- or bisexual, requires the mobilization of educational agents, citizens, and technicians to create a fairer, nondiscriminatory and nonhomophobic society.

Sexology, sexual therapy, and mental health specialists, when sought after, may use appropriate procedures to help nonheterosexuals to live adaptively according to their sexual orientation. If clinical professionals do not feel able to follow international clinical and ethical guidelines, due to personal values or lack of adequate training, it is their duty to those who seek their services, to refer them to clinicians that can conduct psychotherapeutic interventions that are guided by evidence-based intervention models that promote well-being and psychological adjustment (O'Shaughnessy & Speir, 2018; American Psychiatric Association, 2009).

Over the past decade, the positions on sexual orientation in this statement have been reinforced by international and national organizations such as the American Psychological Association (2008), the APA (2018) and the Portuguese Psychological Association (Ordem dos Psicólogos Portugueses, 2017).

The current position is in line with a previous position taken by Portuguese Society of Clinical Sexology 10 years ago.

Acknowledgments

The authors would like to acknowledge the other members of the Board of Directors who are not listed as authors because they were not involved in the writing of the manuscript and who have approved the current statement: Julio Machado Vaz, Graça Santos, Susana Renca, and Sara Ferreira.

ORCID

Patricia M. Pascoal  <http://orcid.org/0000-0002-2810-4034>

Ana Carvalheira  <http://orcid.org/0000-0001-5400-9746>

Jorge Cardoso  <http://orcid.org/0000-0001-6037-3210>

Pedro J. Nobre  <http://orcid.org/0000-0003-2571-6972>

References

- American Psychiatric Association. (2000). *COPP position statement on therapies focused on attempts to change sexual orientation (reparative or conversion therapies)*. Retrieved from: <http://www.psych.org/Departments/EDU/Library/APAOfficialDocumentsandRelated/PositionStatements/200001a.aspx>.
- American Psychiatric Association. (2009). *Resolution on appropriate affirmative responses to sexual orientation distress and change efforts*. Retrieved from <https://www.apa.org/about/policy/sexual-orientation>
- American Psychiatric Association. (2018). *APA reiterates strong opposition to conversion therapy*. Retrieved from: <https://www.psychiatry.org/newsroom/news-releases/apa-reiterates-strong-opposition-to-conversion-therapy>
- American Psychological Association. (2008). *Answers to your questions: For a better understanding of sexual orientation and homosexuality* [PDF file]. Retrieved from www.apa.org/topics/lgbt/orientation.pdf.
- Cochran, S. D., Drescher, J., Kismödi, E., Giami, A., García-Moreno, C., Atalla, E., ... Reed, G. M. (2014). Proposed declassification of disease categories related to sexual orientation in the International Statistical Classification of Diseases and Related Health Problems (ICD-11). *Bulletin of the World Health Organization*, 92(9), 672–679. doi:10.2471/BLT.14.135541
- Drescher, J. (2015). Out of DSM: Depathologizing homosexuality. *Behavioral Sciences*, 5(4), 565–575. doi:10.3390/bs5040565
- Greene, B. (2007). Delivering ethical psychological services to lesbian, gay, and bisexual clients. In K. J. Bieschke, R. M. Perez, & K. A. DeBord (Eds.), *Handbook of counseling and psychotherapy with lesbian, gay, bisexual, and transgender clients* (2nd ed., pp. 181–199). Washington, DC: American Psychological Association. <https://doi.org/10.1037/11482-007>
- Ordem dos Psicólogos Portugueses. (2017). *Guia Orientador da Intervenção Psicológica Com Pessoas Lésbicas, Gays, Bissexuais e Trans* (LGBT) [PDF file]. Lisboa: OPP. Retrieved from: https://www.ordemdospsicologos.pt/ficheiros/documentos/guidelines_opp_lgbt_marco_2017.pdf
- Hall, W. J. (2018). Psychosocial risk and protective factors for depression among lesbian, gay, bisexual, and queer youth: A systematic review. *Journal of Homosexuality*, 65(3), 263–316. doi:10.1080/00918369.2017.1317467
- O'Shaughnessy, T., & Speir, Z. (2018). The state of LGBQ affirmative therapy clinical research: A mixed-methods systematic synthesis. *Psychology of Sexual Orientation and Gender Diversity*, 5(1), 82–98. doi:10.1037/sgd0000259
- Pan American Health Organization and World Health Organization. (2012). *"Therapies" to change sexual orientation lack medical justification and threaten health*. Retrieved from: https://www.paho.org/hq/index.php?option=com_content&view=article&id=6803:2012-therapies-change-sexual-orientation-lack-medical-justification-threaten-health&Itemid=1926&lang=en
- Ross, L. E., Salway, T., Tarasoff, L. A., MacKay, J. M., Hawkins, B. W., & Fehr, C. P. (2018). Prevalence of depression and anxiety among bisexual people compared to gay, lesbian, and heterosexual individuals: A systematic review and meta-analysis. *Journal of Sex Research*, 55(4–5), 435–456. doi:10.1080/00224499.2017.1387755