

## Recent advances in the synthesis of xanthenes and azaxanthenes

Journal:	<i>Organic Chemistry Frontiers</i>
Manuscript ID	Draft
Article Type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Resende, Diana; University of Porto, CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental; University of Porto, Laboratório de Química Orgânica e Farmacêutica, Faculdade de Farmácia Durães, Fernando; University of Porto, CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental; University of Porto, Laboratório de Química Orgânica e Farmacêutica, Faculdade de Farmácia Maia, Miguel; University of Porto, CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental; University of Porto, Laboratório de Química Orgânica e Farmacêutica, Faculdade de Farmácia Sousa, Maria Emília; Universidade do Porto Faculdade de Farmacia, Pinto, Madalena ; University of Porto, LQOF - Laboratório de Química Orgânica e Farmacêutica, Departamento de Ciências Químicas, Faculdade de Farmácia; University of Porto, CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental

## REVIEW

## Recent advances in the synthesis of xanthenes and azaxanthenes

Diana I. S. P. Resende, <sup>a,b,†</sup> Fernando Durães, <sup>a,b,†</sup> Miguel Maia, <sup>a,b,†</sup> Emília Sousa <sup>a,b,\*</sup> and Madalena M. M. Pinto <sup>a,b</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Xanthenes are important *O*-heteroaromatic tricyclic molecules exhibiting a wide range of bioactivities and can represent a showcase to highlight recent techniques and strategies in organic synthesis towards lead discovery and optimization. This review gives an insight of the recent literature disclosed to obtain the xanthone core, including the drug-like azaxanthenes, isosteres of xanthenes, by exploiting the optimization of well-known procedures as well as disruptive developed methodologies. With this review, it is expected to provide a useful chemical toolbox to the synthetic medicinal chemistry focusing in these privileged structures, covering from 2012 to 2020, with some older examples regarding azaxanthenes.

## 1. Introduction

### 2. Synthesis of xanthenes in one reaction step via condensation of two aryl building blocks

#### 2.1 By condensation of a salicylic acid with a phenol derivative

#### 2.2 By condensation of an aryl aldehyde with a phenol derivative

#### 2.3 By condensation of a salicylaldehyde and 1,2-dihaloarenes

#### 2.4 By condensation of a *o*-haloarene-carboxylic acid with arynes

### 3. Synthesis of (aza)xanthenes via benzophenone route

#### 3.1 Synthesis of benzophenones

#### 3.2 Benzophenone cyclization

#### 3.3 Synthesis of azaxanthenes

### 4. Synthesis of (aza)xanthenes via diaryl ether route

#### 4.1 Diaryl ether synthesis

#### 4.2 Cyclization of the diaryl ether

#### 4.3 Synthesis of azaxanthenes

### 5. Synthesis of xanthenes via chromen-4-one derivatives

#### 5.1 From 2-substituted chromen-4-ones

#### 5.2 From 3-substituted chromen-4-ones

#### 5.3 From 2,3-disubstituted chromen-4-ones

### 6. Other methodologies

### 7. Diversifications on the xanthone core

### 8. Conclusions

### 9. Conflicts of interest

### 10. Acknowledgements

### 11. References

## 1. Introduction

Over the past few decades, strategic advances have been made in medicinal chemistry and organic synthesis that accelerated drug discovery. Diversity-oriented synthesis, photo-induced reactions, biorthogonal chemistry with new and eco-friendly reagents, late-stage diversification, and artificial intelligence are among the approaches involved in some of the recent breakthroughs on drug discovery <sup>1, 2</sup>. More than 85% of all biologically-active chemical entities contain a heterocycle which provides a useful tool for lead optimization <sup>3, 4</sup>. This current review aims to emphasize a particular class of heterocycles, the xanthenes, the use of these advances that allow to speeding up their synthesis and leading optimization process.

Xanthenes are naturally occurring compounds that can be found as secondary metabolites in diverse terrestrial and marine plants, fungi, and lichen <sup>5-8</sup>. Chemically, xanthenes (9*H*-xanthen-9-ones) comprise a family of compounds with an oxygen-containing dibenzo- $\gamma$ -pyrone heterocyclic scaffold (Figure 1) <sup>9, 10</sup>. As a privileged structure, this family of compounds is able to provide a wide range of different substitutions and of modulating several biological responses, thus being considered a promising and interesting structure for drug development <sup>11, 12</sup>. Azaxanthenes (Figure 1) are molecules with one or more nitrogen atoms placed in the aromatic moiety of the xanthone chromophore. *N*-heterocycles, present in both natural products and synthetic compounds, constitute important structural motifs in medicinal chemistry, capable of showing a plethora of biological activities <sup>13</sup>. This allied to the privileged *O*-heterocyclic xanthone scaffold, can give rise to new molecules, gifted with interesting activities and drug-likeness, and ultimately to the discovery of new lead compounds <sup>14</sup>. Azaxanthenes also have advantages over acridones, such as being better chromophores and presenting a higher solubility in water <sup>15</sup>.

<sup>a</sup> CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental, Terminal de Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal.

<sup>b</sup> Laboratório de Química Orgânica e Farmacêutica, Faculdade de Farmácia, Universidade do Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal

<sup>†</sup> These authors contributed equally to this work.

\*Corresponding author. E-mail address: esousa@ff.up.pt.




Figure 1. Xanthone and azaxanthones scaffolds.

In terms of structural diversity of the *N*-heterocycle, azaxanthones can be classified into six-membered rings group, if one of the rings is a pyridine or a pyrimidine, or into five-membered rings group, if one of the rings is a pyrazole or a triazole (Figure 1).


In the last decade, both natural and synthetic xanthenes have been reported with diverse biological and pharmacological properties<sup>5, 6, 8, 16</sup>, including antibacterial<sup>8, 17-27</sup>, antifungal<sup>8, 17, 20, 21, 28-30</sup>, antiviral<sup>31</sup>, antioxidant<sup>32-37</sup>, antiobesity<sup>38</sup>, anti-inflammatory<sup>5, 17, 39-41</sup>, anticoagulant<sup>42</sup>, and antitumour<sup>36, 43-58</sup>. Particularly, some xanthenes molecular targets have been disclosed, including clinically important enzymes such as  $\alpha$ -glucosidase<sup>59-63</sup>, topoisomerase<sup>64, 65</sup>, P-glycoprotein<sup>66-68</sup>, and acetylcholinesterase<sup>33, 69, 70</sup> and also protein-protein interactions, such as p53-MDM2<sup>71-74</sup>. Similarly, a variety of biological activities was also disclosed for azaxanthones, depending on the side chains which can be found in their scaffolds. The activities described include antitumour<sup>75, 76</sup>, anti-allergic, bronchodilator, anti-inflammatory, and analeptic<sup>77, 78</sup>. In fact, one azaxanthone, amlexanox, was approved by the Food and Drug Administration (FDA) in 1996, as an antiallergic and anti-inflammatory immunomodulator, used for the treatment of aphthous ulcers<sup>79</sup>. Although its mechanism of action is not fully understood, it is thought to inhibit the release of histamine and leukotrienes from white blood cells<sup>80</sup>. Additionally, amlexanox and derivatives have shown potential as antitumour agents, particularly in tumour cells expressing one or more proteins of the S100 family, which include breast, colon, lung, pancreas, skin, oesophagus, bladder, and other tumours.

The first methods for the synthesis of xanthenes were introduced in by Michael and Kostanecki in 1883-1891<sup>81, 82</sup> and in the 20<sup>th</sup> century and until 2012 several extensive reviews have reported the state of the art regarding the synthesis of xanthenes<sup>9, 10, 83, 84</sup>. More recently reviews have focused on the synthesis of a very specific class of derivatives like carboxyxanthenes<sup>85</sup>, hydroxanthenes<sup>86</sup>, arylxanthenes<sup>87</sup>, and thioxanthenes<sup>88</sup> and, although azaxanthones have demonstrated therapeutic potential and interest as sensitizing chromophores<sup>89-94</sup>, their reports in medicinal chemistry are still sparse in literature. Therefore, in this review the recent exploitation of both scaffolds was considered from a medicinal chemistry point of view with advanced synthetic methodologies applied for obtaining bioactive derivatives being displayed in the sections below.

## 2. Synthesis of xanthenes in one reaction step via condensation of two aryl building blocks

Methods for the synthesis of xanthenes relying in a one-step methodology from readily available building blocks are still quite popular due to their simplicity and possibility of diverse substitution patterns. Four distinct approaches were developed and optimized over the last years (Scheme 1), involving the classical condensation of a salicylic acid with a phenol derivative (2.1), or involving an aryl aldehyde with a phenol derivative (2.2), a salicylaldehyde with 1,2-dihaloarenes (2.3), or a *o*-haloarenecarboxylic acid with arynes (2.4).

## REVIEW




Scheme 1. Methodologies for the synthesis of xanthenes in one reaction step via condensation of two aryl building blocks

### 2.1 Synthesis of xanthenes by condensation of a salicylic acid with a phenol derivative

The classical Grover, Shah and Shah method<sup>95, 96</sup> constitutes a suitable approach for the synthesis of hydroxyxanthenes from easily available salicylic acid derivatives and phenols heated along with zinc chloride in phosphoryl chloride<sup>33, 39, 43, 49, 51, 60, 65, 97-104</sup>. In order to the reaction proceed directly to the desired xanthone, the 2,2'-dihydroxybenzophenone intermediate must contain an additional hydroxyl moiety *ortho* to the carbonyl group. Usually phloroglucinol is used in order to overcome this limitation; however, when other phenol derivatives like resorcinol are used, the synthetic pathway stops at benzophenone form<sup>53, 105</sup>. Consequently, an additional cyclization step is required towards the conversion into xanthenes. These and other limitations over this one-pot strategy towards simple xanthenes led to modifications and development of new synthetic methods<sup>9, 10</sup>. Substitution of phosphorus oxychloride-zinc chloride by phosphorus pentoxide-methanesulfonic acid (Eaton's reagent) as acylation catalyst generally led to better results<sup>18, 19, 23, 24, 33, 35, 40, 50, 54, 59, 63, 70, 106-121</sup>. Since both of these methodologies had previously been extensively reviewed<sup>9, 10</sup>, they will not be discussed herein.

Rare earth metal triflates were used as unique Lewis acids to promote xanthone synthesis catalysed by Yb(OTf)<sub>3</sub> hydrate from 2-substituted benzoic acids and differently substituted phenols under microwave irradiation (Scheme 2)<sup>122</sup>.

Scheme 2. Synthesis of xanthenes via a microwave assisted reaction catalysed by Yb(OTf)<sub>3</sub>

Other metal triflates from the lanthanide series (e.g. La, Eu, and Gd) led to poorest yields, which can be explained by the fact that Yb<sup>3+</sup> is the 'hardest' cation and thus the most oxophilic, due to its smaller ionic radius<sup>122</sup>. Mechanistically, an initial Friedel-Crafts acylation can be considered towards the formation of a benzophenone intermediate, with further ring closure by displacement of the OH group coordinating the metal centre by the OH on the adjacent ring. Combination of salicylic, gentistic or 5-nitrosalicylic acids with *m*-cresol, orcinol, phloroglucinol, resorcinol, pyrogallol, *p*-nitrophenol, *p*-chlorophenol, 7-hydroxycoumarin or 4-methyl-7-hydroxycoumarin led to the corresponding xanthenes **3** in high yields (19 examples, 72-98%)<sup>122</sup>.


### 2.2 Synthesis of xanthenes by condensation of an aryl aldehyde with a phenol derivative

Highly selective and efficient transition metals like Cu and Fe catalytic systems have been recently employed for the synthesis of xanthenes due to the associated concept of eco-friendliness, considering atom economy, low toxicity and atom efficiency<sup>123-126</sup>. In the scope of a study on *ortho*-acylation of phenols with various aryl aldehydes using CuCl<sub>2</sub> as the catalyst in the presence of triphenylphosphine, it was found that benzophenones were obtained when benzaldehyde or 3- or 4-substituted aryl aldehydes were employed. Nevertheless, when 2-substituted aryl aldehydes reacted with phenols, xanthenes were obtained (Table 1, Method A), first *via ortho*-acylation of phenols with 2-substituted aryl aldehydes with further ring closure under basic conditions, with *ortho*-substituents of the aryl aldehydes serving as leaving groups (LG)<sup>123</sup>. The scope of the reaction of 2-nitrobenzaldehydes and phenols was investigated after an initial screening regarding leaving groups, showing that electron-donating groups (EDG) (alkoxy, alkyl, aryl, and halides) were well tolerated on phenols, promoting the reaction. On the other hand, electron-withdrawing groups (EWG) (NO<sub>2</sub> or CN) block the reaction completely, being the main drawback of this methodology<sup>123</sup>. Although these results are promising for eco-friendly reasons, they still employ ligands and expensive catalysts which are not recycled. To surpass those disadvantages, Menendéz et al.<sup>124</sup> optimized the reaction

REVIEW

Organic Chemistry Frontiers

**Table 1.** Synthesis of xanthenes via a one-step ortho-acylation of phenols with 2-substituted aldehydes



**Method A,** Y = NO<sub>2</sub>

CuCl<sub>2</sub>, PPh<sub>3</sub>, K<sub>3</sub>PO<sub>4</sub>, toluene, 110 °C, air, 24 h

**Method C,** Y = NO<sub>2</sub>, Cl, Br

Fe<sub>3</sub>O<sub>4</sub> NPs, K<sub>2</sub>CO<sub>3</sub>, toluene, 110 °C, air, 24h

**Method B,** Y = NO<sub>2</sub>

CuNPs, K<sub>3</sub>PO<sub>4</sub>, toluene, Ar, 2 h

**Method D,** Y = NO<sub>2</sub>

nano-CuFe<sub>2</sub>O<sub>4</sub>, K<sub>3</sub>PO<sub>4</sub>, DMF, 80 °C, air


Entry	Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	% Yield/ Ref.			
							Method A <sup>123,[a]</sup>	Method B <sup>124,[a]</sup>	Method C <sup>125,[b]</sup>	Method D <sup>126,[b]</sup>
1	NO <sub>2</sub>	H	H	H	OCH <sub>3</sub>	H	87 (74)	98 (89)	-	-
2	NO <sub>2</sub>	H	H	H	<i>i</i> Pr	H	92 (81)	-	-	-
3	NO <sub>2</sub>	H	H	H	H	H	81 (70)	84 (76)	65	89
4	NO <sub>2</sub>	H	H	H	I	H	77 (62)	-	-	80
5	NO <sub>2</sub>	H	I	H	H	H	73 (62)	-	-	78
6	NO <sub>2</sub>	H	H	H	Br	H	72 (59)	78 (70)	-	86
7	NO <sub>2</sub>	H	H	H	Cl	H	82 (73)	-	-	82
8	NO <sub>2</sub>	H	Cl	H	H	H	76 (60)	-	-	74
9	NO <sub>2</sub>	H	H	F	H	H	70 (58)	-	-	-
10	NO <sub>2</sub>	H	H	H	F	H	71 (55)	-	-	79
11	NO <sub>2</sub>	H	Ph	H	H	H	69 (55)	-	-	-
12	NO <sub>2</sub>	H	C <sub>4</sub> H <sub>4</sub>	H	H	43 (30)	85 (77)	60	79	-
13	NO <sub>2</sub>	H	Cl	H	Cl	H	75 (64)	-	-	-
14	NO <sub>2</sub>	H	H	CH <sub>3</sub>	CH <sub>3</sub>	H	84 (72)	-	-	80
15	NO <sub>2</sub>	H	CH <sub>3</sub>	H	CH <sub>3</sub>	H	90 (80)	-	-	-
16	NO <sub>2</sub>	H	H	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	73 (61)	-	-	-
17	NO <sub>2</sub>	OCH <sub>3</sub>	H	H	OCH <sub>3</sub>	H	83 (70)	85 (72)	-	-
18	NO <sub>2</sub>	OCH <sub>3</sub>	H	H	<i>i</i> Pr	H	82 (70)	-	-	-
19	NO <sub>2</sub>	OCH <sub>3</sub>	H	H	Cl	H	80 (69)	-	-	-
20	NO <sub>2</sub>	OCH <sub>3</sub>	Cl	H	Cl	H	68 (57)	-	-	-
21	NO <sub>2</sub>	H	H	OCH <sub>3</sub>	H	H	-	97 (86)	-	-
22	NO <sub>2</sub>	H	OCH <sub>3</sub>	H	H	H	-	91 (84)	-	-
23	NO <sub>2</sub>	H	H	H	NO <sub>2</sub>	H	-	60 (51)	-	-
24	NO <sub>2</sub>	H	H	H	CF <sub>3</sub>	H	-	63 (54)	-	-
25	NO <sub>2</sub>	H	CH <sub>3</sub>	H	H	H	-	89 (80)	-	76
26	NO <sub>2</sub>	H	H	OH	H	H	-	75 (68)	-	-
27	Cl	H	H	H	H	H	-	-	55	-
28	Br	H	H	H	H	H	-	-	50	-
29	NO <sub>2</sub>	H	H	H	<i>t</i> Bu	H	-	-	72	84
30	Br	H	H	H	<i>t</i> Bu	H	-	-	55	-
31	NO <sub>2</sub>	H	H	CH <sub>3</sub>	H	H	-	-	-	92
32	NO <sub>2</sub>	H	H	H	CH <sub>3</sub>	H	-	-	-	85
33	NO <sub>2</sub>	H	H	C <sub>4</sub> H <sub>4</sub>	H	-	-	-	80	-
34	NO <sub>2</sub>	CH <sub>3</sub> <sup>[c]</sup>	H	H	H	H	-	-	-	78
35	NO <sub>2</sub>	Br	H	H	H	H	-	-	-	75
36	NO <sub>2</sub>	Cl	H	H	H	H	-	-	-	70

[a] <sup>1</sup>H NMR yield (isolated yield); [b] isolated yields; [c] 4-methyl-2-nitrobenzaldehyde

conditions using a magnetic nanocatalyst consisting of CuNPs (NP = nanoparticles) on silica coated maghemite to catalyze the reaction between *ortho*-substituted benzaldehydes and phenols (Table 1, Method B). Provided with a high surface area, nano catalysts were used to afford a more significant advancement in catalytic activity over conventional catalytic methods. This synthetic protocol was compatible with the presence of a variety of functional groups in the starting phenolic building blocks, affording the desired xanthenes in good to excellent yields, with recovery of the catalyst<sup>124</sup>. Ramani<sup>125</sup> used Fe<sub>3</sub>O<sub>4</sub> nanoparticles under ligand-free conditions to catalyze *ortho*-benzoylation of phenols with aryl aldehydes for the synthesis of xanthenes, with lower yields than the reported for the latter two methodologies (Table 1, Method C). Later, Venkanna<sup>126</sup> used magnetically separable nano-CuFe<sub>2</sub>O<sub>4</sub> to catalyze *ortho*-acylation (Table 1, Method D). Xanthone derivatives were generally obtained in good yields from the condensation of diverse phenols with 2-nitrobenzaldehydes using nano-CuFe<sub>2</sub>O<sub>4</sub> as catalyst, K<sub>3</sub>PO<sub>4</sub> and DMF at 80 °C, in an open flask, for 10 h. Comparing the four methodologies (Table 1), and looking at characteristics of eco-friendliness, reaction conditions and yields, methods B and D seem to be more suitable for the synthesis of xanthenes from 2-nitrobenzaldehydes and phenols, with a good tolerance for different substituent groups on the phenol moiety.

### 2.3 Synthesis of xanthenes by condensation of a salicylaldehyde and 1,2-dihaloarenes

Aside *ortho*-acylation of phenols with 2-substituted aldehydes, the xanthone core can also be constructed through a selective palladium-catalysed acylation/nucleophilic aromatic substitution (S<sub>N</sub>Ar) approach of salicylaldehyde derivatives with aryl halides (Scheme 3)<sup>127, 128</sup>. The first methodology reported consisted in the use of a bulky electron-rich trialkylphosphine ligand, *n*-BuPAd<sub>2</sub>, which prevents the decomposition of the palladium catalyst, observed with the use of other ligands<sup>128</sup>. The scope of the reaction was extended to 22 different xanthenes (41-81% yield); however, only activated salicylaldehydes were tolerated, long reaction times (12 h) were needed, and it was impractical to recover and reuse the catalyst. A rapid and convenient ligand-free alternative was later reported for this direct annulation using a recoverable



Scheme 3. Synthesis of xanthenes via palladium-catalysed acylation/S<sub>N</sub>Ar approach

palladium nanocatalyst supported on a green biochar under microwave irradiation, with a drastic reduction on reaction times<sup>127</sup>. This eco-friendly strategy proceeded in very good yields (54-88%) and with high regioselectivity for a scope of 20 xanthone derivatives.

### 2.4 Synthesis of xanthenes by condensation of *o*-haloarene-carboxylic acids with arynes


Dubrovskiy and Larock<sup>129</sup> reported the synthesis of xanthenes via an intermolecular C-O addition of *o*-halobenzoic acids **10** to arynes **11** employing CsF and THF at 125 °C for 24 h. Albeit *o*-chloro and *o*-iodo benzoic acids provided poor yields, *o*-fluorobenzoic acid afforded the desired xanthenes **12** in good yields (Scheme 4). Diversification of substituents either in the benzoic acid building block or in the arylene moiety was well tolerated, and nine different xanthone derivatives were obtained, including an azaxanthone.

## 3. Synthesis of xanthenes via benzophenone route

The limitations of some of the earlier one step methodologies (e.g. Grover, Shah and Shah, 2.1) led to the development of several multi-step approaches. The synthesis of xanthenes via benzophenone route is one of the most used approaches in which the benzophenone derivatives are easily accessed through (poly)oxygenated aromatic compounds with a benzoic acid derivative (3.1), followed by the cyclization of the benzophenone (3.2) into the xanthone scaffold (Scheme 5), usually achieved by base-catalysed intramolecular nucleophilic aromatic substitution. Considering the high versatility of this route, examples on the total synthesis of natural products using this strategy in the last decade<sup>130-136</sup> are represented in Figure 2.

### 3.1 Synthesis of benzophenones

Friedel-Crafts acylation continues to be the method of choice for the synthesis of the benzophenone moiety, when xanthenes are obtained via benzophenone route (3.1.1)<sup>28, 47, 66, 71, 137-140</sup>. Additional approaches for the synthesis of the benzophenone moiety include the halogen-lithium exchange (3.1.2) and NHC-catalysed arylation (3.1.3) (Scheme 6).




Scheme 4. Synthesis of xanthenes via an intermolecular C-O addition of *o*-halobenzoic acids to arynes



REVIEW

Organic Chemistry Frontiers



Scheme 5. Strategies for the synthesis of xanthenes via benzophenone

Other methodologies were reported for the synthesis of benzophenones via a *ortho*-acylation of phenols using copper<sup>123</sup> and iron-mediated<sup>125</sup> approaches, or via a reductive decarbonylative coupling<sup>141</sup>. However, further investigations led to the attainment of the corresponding xanthone in one step and, therefore, they are addressed in the appropriate sections (2.2 and 7.1).

**3.1.1 Synthesis of benzophenones via Friedel-Crafts acylation of (poly)oxygenated aromatic compounds with a benzoic acid derivative.** Over the last years, several examples on the synthesis of benzophenones via Friedel-Crafts acylation have been reported, mainly via classical reaction conditions employing an acid chloride and  $\text{AlCl}_3$  as Lewis acid catalyst (Scheme 6A)<sup>28, 47, 66, 71, 137-139</sup>. When two methoxy groups are present in both  $R^1$  and  $R^9$  three different benzophenone derivatives can be formed: a benzophenone methoxylated at  $R^1$  and  $R^9$ , and two benzophenone derivatives which result from the demethylation of the methoxy group in  $R^1$  or  $R^9$  (position *peri*-carbonyl). Hence, to avoid losses, usually the benzophenone derivatives are not isolated and the reaction crude is used directly towards cyclization to xanthenes<sup>28, 66, 71, 138</sup>. Other interesting strategies have been developed in the last years for the synthesis of benzophenones, being worth to

highlight one in which the carbonyl bridge is introduced via a mild Friedel Crafts reaction using trifluoroacetic anhydride (TFAA) (Scheme 6A)<sup>140</sup>. This methodology was developed in the scope of a study to find novel PGAM1 inhibitors and involves the in-situ formation of a mixed anhydride that results from the reaction of the benzoic acid derivative with TFAA, in order to make the carbonyl adjacent to the aromatic ring more susceptible to nucleophilic attack<sup>140</sup>. Due to limitations on the tolerated substituents that can be present at either the benzoic acid derivative or the (poly)oxygenated aromatic compound in order for cyclization to occur, this methodology is usually employed as starting point for the synthesis of more complex xanthone derivatives.

**3.1.2 Synthesis of benzophenones via a halogen–lithium exchange.** The halogen–lithium exchange benzophenone cyclization is not used as commonly as the Friedel-Crafts acylation, but can still be considered a valuable tool for the synthesis of benzophenone derivatives. Studies on the synthesis of 12-hydroxy-2,2-dimethyl-3,4-dihydropyran[3,2*b*]xanthene-6(2*H*)-one (**19**), a p53-MDM2 inhibitor<sup>74</sup>, led to benzophenone **18**, obtained from halogen–lithium exchange of previously synthesized benzopyran **16** and *O*-protected MOM (methoxymethyl) benzyl salicylate **17** (Scheme 6B)<sup>46</sup>.





Figure 2. Recent examples on the total synthesis of natural products using benzophenone route

## A. Synthesis of benzophenones via Friedel-Crafts acylation




## Reaction conditions

- a)  $\text{AlCl}_3$ ,  $\text{Et}_2\text{O}$ , rt, 5–18 h, 62–65%, when isolated  
 b)  $\text{AlCl}_3$ ,  $\text{Et}_2\text{O}$ , 0–10 °C, yield not determined  
 c) TFAA,  $\text{CH}_2\text{Cl}_2$ , 0 °C, 66% yield


Y = Cl or OH

 $\text{R}^1 = \text{R}^7 = \text{OCH}_3$  $\text{R}^2 = \text{R}^3 = \text{R}^4 = \text{R}^6 = \text{H}$  or  $\text{OCH}_3$  $\text{R}^5 = \text{R}^8 = \text{H}$ ,  $\text{CH}_3$  or  $\text{OCH}_3$  $\text{R}^8 = \text{OCH}_3$  or  $\text{OBn}$ 

## B. Synthesis of benzophenones via a halogen-lithium exchange



## C. Synthesis of benzophenones via a NHC-catalyzed arylation



Scheme 6. Synthesis of benzophenones via Friedel-Crafts acylation of (poly)oxygenated aromatic compounds with a benzoic acid derivative

**3.1.3 Synthesis of benzophenones via a NHC-catalysed arylation.** *N*-heterocyclic carbenes (NHC) can be employed both as ligands for transition-metal systems or as small-molecule organocatalysts for chemical synthesis. In a nucleophilic arylation catalysed by NHC **22**, the 4-fluoro group of trifluoronitrobenzene (**20**) is replaced by the aroyl group originating from aldehyde **21** to afford the corresponding benzophenone **23** (Scheme 6C) <sup>133</sup>. This versatile methodology was applied in the total synthesis of termicalcolanone A (Figure 2), an anticancer natural xanthone <sup>133</sup>.

## 3.2 Benzophenone cyclization

The base-catalysed intramolecular nucleophilic aromatic substitution is undoubtedly the most frequently used strategy for the cyclization of benzophenones to xanthenes (3.2.1). This methodology involves the presence of a hydroxyl group in position 2 and a functional group, normally a methoxy, which can behave as a leaving group in position 2' of the benzophenone, usually in the form of methanol. Other methodologies include oxidative cyclization (3.2.2) with new and greener reaction conditions being reported in the last years.

## 3.2.1 Cyclization of benzophenones to xanthenes via a base-catalysed intramolecular nucleophilic aromatic substitution.

Although the classical methodologies are widely used <sup>130, 132, 137–140</sup>, some improvements, especially in terms of yields and shorter reaction times, were achieved with microwave-assisted organic synthesis (MAOS) <sup>28, 46, 47, 66, 71</sup>. Common bases used in this strategy are NaOH <sup>28, 47, 66, 71, 138</sup>, tetramethylammonium hydroxide (TMAH) <sup>137, 140</sup>, and  $\text{K}_2\text{CO}_3$  <sup>132, 139</sup>. A greener approach was also reported using a combination of base/ $\text{H}_2\text{O}$  <sup>142</sup> (Scheme 7A).

In the presence of 2-halo-2'-hydroxybenzophenones, a transition-metal-free intramolecular Ullmann-type *O*-arylation reaction (Scheme 7B) can be employed for the synthesis of the corresponding xanthenes **27**, either using  $\text{K}_2\text{CO}_3$  or tetrabutylammonium hydroxide (TBAOH) as base <sup>143</sup>. The latter is an efficient, rapid, and green method which uses microwave irradiation, resulting in short reaction time and excellent yields. *O*-Arylation reaction proceeds smoothly, regardless of the nature of the halogen leaving group present in the aromatic moiety, with several functional groups being well-tolerated in all the examples, and extendable to the synthesis of




## REVIEW

## Organic Chemistry Frontiers

azaxanthenes. Despite the desired xanthenes **27** were produced in comparatively lower yields when in the presence of


### A. Synthesis of xanthenes via a base-catalyzed intramolecular nucleophilic aromatic substitution



#### Experimental conditions

- TMAH, pyridine, H<sub>2</sub>O, reflux, 36 h, 2 examples, 83-92% yield
- NaOH, CH<sub>3</sub>OH/H<sub>2</sub>O, 110 °C, 48 h, 2 examples, 58-90% yield
- NaOH, CH<sub>3</sub>OH/H<sub>2</sub>O, MW, 30 min, 1 example, 65% yield
- K<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>OH, H<sub>2</sub>O, 28 h, 2 examples, n.d. yield
- NaOH, CH<sub>3</sub>OH/H<sub>2</sub>O, MW, 100 °C, 6h, 2 examples, 15-63% yield (2 steps)


### B. Synthesis of xanthenes and azaxanthenes via intramolecular Ullmann-type O-arylation



#### Experimental conditions

- K<sub>2</sub>CO<sub>3</sub>, acetone, 25 °C, 2 h, 1 example, 96% yield
- K<sub>2</sub>CO<sub>3</sub>, DMF, 75 °C, 1 example, 63% yield (3 steps)
- TBAOH, H<sub>2</sub>O, 120 °C, MW, 4 min, 20 examples, 98-99% yield

### C. Cyclization of benzophenones to xanthenes via an oxidative cyclization



#### Experimental conditions

- Ag<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN, 100 °C, 2 h, 1 example, 62% yield
- CAS, H<sub>2</sub>O/CHCl<sub>3</sub>/CH<sub>3</sub>CN, rt, 24 h, 4 examples, 21-91% yield
- CAS, H<sub>2</sub>O/CHCl<sub>3</sub>/CH<sub>3</sub>CN, rt, 18 h, 1 example, 60% yield

Scheme 7. Synthesis of xanthenes via benzophenone cyclization

1-hydroxynaphthalene, in 4 min under MW irradiation, quantitative yields could be obtained when the reaction time was extended to 8 min. Azaxanthone was prepared in two minutes with an excellent yield (98%)<sup>143</sup>.

**3.2.2 Cyclization of benzophenones to xanthenes via an oxidative cyclization.** Despite not being used as often as the base-catalysed intramolecular nucleophilic aromatic substitution, methodologies based on the oxidative cyclization of the benzophenone are also employed for the synthesis of xanthone derivatives. Studies on the cyclization of the readily available model substrate 2,3',5-trihydroxy benzophenone **28** revealed that the combination of Ag<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN, at 100 °C for 2 h gave the best results towards the cyclization product, with the hydroxyl group being crucial for cyclization that occurs via a single-electron transfer mechanism<sup>144</sup> (Scheme 7C). To obtain 1,5,7-trisubstituted-3-pyridylxanthenes with potential as insecticides, the target precursor 1,5,7-trihydroxy-3-chloroxanthone was cyclized using the optimized

reaction conditions, in the route for the synthesis of several 1,5,7-trisubstituted-3-pyridylxanthenes<sup>144</sup>. Ceric ammonium sulfate (CAS)<sup>145, 146</sup> and ceric ammonium nitrate (CAN)<sup>146</sup>-mediated oxidative cyclization of benzophenones was also reported, although some limitations were observed with the formation of a xanthone-related dione. A possible mechanism for the formation of this secondary product was proposed with the formation of a radical cation with subsequent nucleophilic addition of the phenol, oxidation with CAN, and addition of water followed by elimination of methanol<sup>145</sup>.

### 3.3 Synthesis of azaxanthenes


The synthesis of azaxanthenes via the benzophenone route usually employs the preparation of the required benzophenone through Friedel-Crafts acylation (Scheme 8A), deprotocuration-arylation (Scheme 8B), or even the formation of an intermediate alcohol via metal-mediated strategies, which is further oxidized to the corresponding

ketone (Scheme 8C and 8D). Cyclization is typically achieved via  $S_NAr$ .


**3.3.1 Synthesis of azaxanthenes through reaction between 2-chloronicotinic acid and phloroglucinol.** Kolokythas *et al.*<sup>76</sup> have studied the potential of pyranoxanthenones and pyranothioxanthenones, as potential cytotoxic agents closely related

to the broad spectrum antitumour alkaloid acronycine (Scheme 8A). As an outcome for this work, the carbocyclic moiety was replaced by the pyridine heterocycle isostere in the xanthone chromophore. Similarly to the work of Villani *et al.*<sup>78</sup>, formation of the corresponding acyl chloride of 2-chloronicotinic acid (**30**) and further reaction with phloroglucinol **31**, provided ketone **32** which was then cyclized to dihydroazaxanthone **33** in 51% yield<sup>76</sup>.


#### A. Synthesis of azaxanthenes through reaction between 2-chloronicotinic acid and phloroglucinol




#### B. Synthesis of azaxanthenes by deprotocupration-arylation followed by intramolecular direct arylation



#### C. Synthesis of 2-azaxanthone using a lithiation strategy



#### D. Synthesis of 4-azaxanthone by trans metalation



## REVIEW

Scheme 8. Synthesis of azaxanthenes via benzophenone route

### 3.3.2 Synthesis of azaxanthenes through deprotocupration-arylation followed by intramolecular direct arylation.

Lithiocuprates, prepared from copper(II) chloride and lithium 2,2,6,6-tetramethylpiperidide (LTMP) can be used as bases for the deprotonative metalation of aromatic compounds. By using palladium as a catalyst for ring closure, it is possible to obtain not only the expected azafluorenones, but also azaxanthenes, through cyclization, in the presence of oxygen-containing substituents or reagents<sup>147</sup>. Synthesis of azaxanthenes **42–44** was achieved by using this approach. 4,5-Diazaxanthone (**42**), was prepared from 2-chloropyridine (**34**) and LTMP, which stirred for two hours at 0 °C in THF, before the addition of the corresponding aryl chloride (Scheme 8B). The 4,5-diazaxanthone **42** was produced in 45% overall yield (2 steps) after treatment with palladium(II) acetate and trichlorohexylphosphine tetrafluoroborate. The second azaxanthone, 4-azaxanthone (**43**), was synthesized similarly, starting from 2-methoxypyridine (**35**) and chlorobenzyl chloride in 8% overall yield (2 steps). The second step was slightly different, and although Pd(OAc)<sub>2</sub> was used as a catalyst, tri-*tert*-butylphosphonium tetrafluoroborate was also employed (Scheme 8B). 2-Azaxanthone (**44**), was obtained through the same method as 4,5-diazaxanthone (**42**), using 4-methoxypyridine (**36**) and chlorobenzyl chloride (Scheme 8B), yielding 29% of the desired compound (overall yield for 2 steps)<sup>147</sup>.

### 3.3.3 Synthesis of azaxanthenes through metalation – lithiation.

Metalation of  $\pi$ -deficient heterocycles, such as pyridine, has proven to be a good strategy towards obtaining compounds that could serve as starting materials for azaxanthenes. This methodology can be a powerful functionalization approach, since electrophilic substitutions are frequently hard to achieve in such molecules. Based on the fact that halogens can induce *ortho*-lithiation of pyridine, Marsais *et al.*<sup>148</sup> explored this route and one of their experiments led to the synthesis of 2-azaxanthone (**44**). The metalation of the starting material, 4-fluoropyridine (**45**), was achieved with complete chemoselectivity using lithium diisopropylamide (LDA) and *o*-anisaldehyde (**46**). The formation of butylated side products when using butyllithium attested the advantages of using LDA as lithiation agent. The fact that the product, (4-fluoro-3-pyridyl)-2-methoxyphenylmethanol (**47**), precipitated with diethyl ether also constituted an advantage, as higher metalation yields could be reached. Oxidation of **47** with manganese dioxide, led to (4-fluoro-3-pyridyl)-2-methoxyphenylmethanone (**48**) which, when heated with pyridinium chloride yielded 2-azaxanthone (**44**), in 49% overall yield, obtained over 3 steps (Scheme 8C1)<sup>148</sup>. A similar approach was used for the synthesis of halo-azaxanthenes, starting from 2,6-difluoropyridine (**49**) and 5-bromo-2-methoxybenzaldehyde (**50**). A *tert*-butyldimethyl silyl (TBS) ether protection of the formed hydroxyl moiety and installation of a Bu<sub>3</sub>Sn group after the second fluorine-directed *ortho*-lithiation of the pyridine were used as key steps in the synthesis of trisubstituted azaxanthone **55** in 35% yield (overall yield considering 7 steps) (Scheme 8C2)<sup>149</sup>.

**3.3.4 Synthesis of azaxanthenes through transmetalation.** It is clear that metalation reactions are useful, with direct reactions with electrophiles or transmetalation, leading to cross-coupling reactions, being the most important. However, the lithiation reactions previously described require low temperatures, and, added to the fact that halogen-lithium exchange is extremely temperature-sensitive, that can be limiting, making them difficult to perform in an industrial setting<sup>150</sup>. Therefore, the search for alternative reactions led to the development of pyridylmagnesium reagents that could either take part in electrophilic trapping or coupling reactions. In this scope, halogen-magnesium exchange methodology was preferred to the Grignard reaction, as the direct access by oxidative addition of magnesium to the halopyridine is challenging to accomplish, even with magnesium activation<sup>150</sup>. Studies on synthetic routes leading to new pyridine derivatives through bromine-magnesium exchange<sup>148, 150</sup> revealed that isopropylmagnesium chloride (*i*PrMgCl) was the best exchange reagent, and THF the best solvent. The fact that this reaction occurs at room temperature posed as an advantage. 4-Azaxanthone (**37**) was prepared starting from 3-bromo-2-chloropyridine (**51**) (Scheme 8D). Metalation with *i*PrMgCl, using THF as solvent, at room temperature and further reaction with *o*-anisaldehyde (**41**) resulted in the formation of alcohol (**52**), which was then subject to oxidation and ring closure reactions affording **37** as described before, in 55% overall yield (3 steps).

## 4. Synthesis of xanthenes via diaryl ether route

The via diaryl ether synthesis comprises the reaction between aryl halides and phenols which are further cyclized to the xanthone core. Like the benzophenone route, the diaryl ether route can then be divided in two steps: the synthesis of the diaryl ether (4.1) and the cyclization of the diaryl ether (4.2) (Scheme 9). This approach is particularly useful in the synthesis of 1,2-dioxygenated xanthenes, when the pattern of substitution of the target xanthone cannot be obtained following the electrophilic aromatic substitution rules and/or is also dependent on the presence of bulky groups giving rise to isomeric mixtures. Nevertheless, due to generally higher yields obtained with intermolecular acylations when compared to Ullmann ether syntheses, the recent examples on the total synthesis of natural products using diaryl ether route are quite scarce, contrary to the existing ones via benzophenone route (Figure 3).

### 4.1 Diaryl ether synthesis

With the extensive development of several methodologies towards xanthone synthesis, either through the benzophenone route (section 3) or even by one-step methodologies (section 2) or newer strategies (section 7), fewer examples involving other





Figure 3. Recent examples on the total synthesis of natural products using diaryl ether route



Scheme 9. Strategies for the synthesis of xanthenes via diaryl ether

procedures have been described in recent times. When used, diaryl ether synthesis is usually achieved via the classic Ullmann condensation (4.1.1), which consists in the coupling of an aryl halide with a phenol (Scheme 4).

**4.1.1 Synthesis of diaryl ether by coupling of an aryl halide with a phenol - Ullmann condensation.** While some modifications on the coupling of an aryl halide with a phenol using Ullmann conditions have been described in the last years, copper continues to be the metal of choice for this coupling. Common bases for Ullmann type reactions are  $\text{K}_2\text{CO}_3$  and  $\text{Cs}_2\text{CO}_3$  and some variations in the copper catalyst include the use of  $\text{CuI}$ <sup>45, 151</sup>,  $\text{CuO}$ <sup>152</sup>,  $\text{CuCl}$ <sup>153</sup> or  $\text{Cu/Cu}_2\text{O}$ <sup>154</sup>. Since this methodology has already been extensively reviewed, recent examples will be summarized in Scheme 10 and will not be widely discussed. However, it is worth to mention a new copper-(I)-catalysed approach that, when compared to the classical Ullmann reaction protocols requires only small amounts of copper catalyst and mild temperatures, with better overall yields, even when tested at a larger scale. The Buchwald's  $\text{Cu}(\text{OTf})_2$ -benzene complex proved to be extremely effective against the classic  $\text{CuI}$  when combined with 4-dimethylaminopyridine (4-DMAP), to give the desired diaryl ether in high yields<sup>155</sup>. One of the synthesized xanthenes showed typical xanthone fluorescence properties suitable for studying folding processes of polypeptides by triplet–triplet energy transfer (TTET)<sup>155</sup>.

## 4.2 Cyclization of the diaryl ether

To be used as substrates in the synthesis of xanthenes, diaryl ethers should contain certain molecular characteristics in order to the next step, cyclization, can occur. Over the last years, several approaches to the xanthone skeleton from a range of functionalized diaryl ethers via varied mechanisms have been developed, depending largely on the functionality existing in


*ortho* position to the ether (Scheme 9,  $Y$ ). When in the presence of an aldehyde derivative ( $Y = \text{CHO}$ ), the most common methodology is via cross dehydrogenative coupling (CDC, 4.2.1). This strategy has received increasingly attention in recent years due to the fact that it does not require pre-functionalized coupling partners, can be extended to a wide scope of derivatives, involves few steps, and consequently high atom economy. In the presence of a carboxylic acid derivative ( $Y = \text{COOH}$ ,  $\text{COCl}$ ,  $\text{COOBn}$ ), three approaches can be followed, depending on the nature of the derivative: an acyl radical cyclization via photoredox catalysis using methylene blue as a photosensitizer (4.2.2,  $Y = \text{COOH}$ ), a metal-free variant of the intramolecular Friedel–Crafts acylation (4.2.3,  $Y = \text{COCl}$ ), and a  $\text{FeCl}_3$  mediated direct intramolecular acylation of esters (4.2.4,  $Y = \text{COOBn}$ ). However, the presence of a carboxylic acid derivative is not mandatory, and other substrates, as 2-iodo diaryl ethers ( $Y = \text{I}$ ) or the corresponding diazonium salts ( $Y = \text{N}_2^+ \text{BF}_4^-$ ) can be used for the synthesis of xanthenes in a palladium-catalysed carbonylation/ $\text{C-H}$  activation (4.2.5). Other kind of cyclization strategies have also been reported in which the xanthone core can be obtained through a  $\text{Co}^{\text{I}}$ -catalysed Barbier reaction of aromatic halides with aromatic aldehydes and imines<sup>156</sup>, a  $\text{PhI}(\text{OAc})_2\text{-BF}_3\text{-OEt}_2$  mediated domino imine activation, intramolecular  $\text{C-C}$  bond formation and  $\beta$ -elimination<sup>157</sup>, or a quaternary ammonium salt-promoted intramolecular dehydrogenative arylation of aldehydes<sup>158</sup>.

### 4.2.1 Cyclization of diaryl ethers to xanthenes by a cross dehydrogenative coupling.

After the pioneer work of Wang et

REVIEW

Organic Chemistry Frontiers




Experimental conditions

- CuI, Cs<sub>2</sub>CO<sub>3</sub>, *N,N*-dimethyl glycine, dioxane, N<sub>2</sub>, 90 °C, 14 h, 54% yield
- CuI, Cs<sub>2</sub>CO<sub>3</sub>, *N,N*-dimethyl glycine, dioxane, N<sub>2</sub>, 90 °C, 18 h, 26% yield
- CuO, K<sub>2</sub>CO<sub>3</sub>, pyridine, reflux, 24 h, 62% yield
- CuCl, Cs<sub>2</sub>CO<sub>3</sub>, TDA-1, dioxane, reflux, N<sub>2</sub>, overnight, yield not determined
- Cu, Na<sub>2</sub>CO<sub>3</sub>, pyridine, reflux, N<sub>2</sub>, 26 h, 39-42% yield (2 examples)
- (CuOTf)<sub>2</sub>.PhH, 4-DMAP, Cs<sub>2</sub>CO<sub>3</sub>, dioxane, reflux, 36 h, 91% yield

Scheme 10. Synthesis of diaryl ether by Ullmann condensation

al. <sup>159</sup>, in 2012 on the synthesis of xanthenes via a rhodium-catalysed cross dehydrogenative coupling (CDC) starting from 2-aryloxybenzaldehydes, alternative transition-metal catalysed strategies to construct the xanthone scaffold have been developed (Scheme 11) <sup>160-163</sup>. For example, Wertz et al. <sup>160</sup> reported an elegant and straightforward procedure for the synthesis of the xanthone core through a FeCp<sub>2</sub>-catalysed CDC reaction via a base-promoted homolytic aromatic substitution. Under optimized conditions, several 2-aryloxybenzaldehydes bearing both electron-donating and electron-withdrawing substituents were coupled efficiently to give the corresponding xanthenes with yields ranging from 30 to 78% (Scheme 11). Later, Manna et al. <sup>161</sup> reported an *ortho* C–H bond functionalization involving dichloro(*p*-cymene)ruthenium(II) dimer [[RuCl<sub>2</sub>(*p*-cymene)<sub>2</sub>]<sub>2</sub> catalysed C–C bond formation. When compared to other metal-mediated CDC methodologies, this protocol does not require any expensive metal-catalyst and uses a commercially available cheap oxidant. The authors extended the scope of this reaction not only to xanthenes, but also to thioxanthenes and fluorenones in moderate to good yields <sup>161</sup>. Recently, Bao et al. <sup>162</sup> described a Cu(0)/Selectfluor system-catalysed intramolecular Csp<sup>2</sup>-H/Csp<sup>2</sup>-H bond CDC of 2-aryloxybenzaldehydes (Scheme 11).

Interestingly, when in the presence of a substituent in *ortho* position to the formyl group on the 2-aryloxybenzaldehyde, regioisomers were obtained. The presence of substituents at the *ortho*, *meta*, or *para* positions relatively to the formyl group are well tolerated, being the products obtained in good yields, except in the case of OPh, affording the desired product in lower yield (49%). Besides clear advantages related to the use of an inexpensive and low-toxic copper catalyst and relative mild reaction conditions, this procedure also accepts a wide variety of substituent groups such as F, Cl, Br, OCF<sub>3</sub>, CF<sub>3</sub>, Ph, OPh, *i*Pr, *i*Bu, which provides opportunities for further molecular modifications. The authors were able to expand the use of this methodology for the synthesis of 9*H*-thioxanthene-9-one and 10,10-dioxide and phenanthridin-6(5*H*)-one derivatives <sup>162</sup> and, interestingly, this process could be conducted successfully on a gram scale for a simple oxygenated xanthone (80% yield). A plausible mechanism for the [RuCl<sub>2</sub>(*p*-cymene)<sub>2</sub>]-catalysed cyclization reaction was proposed and involves the formation of



Metal-mediated CDC

- RhCl<sub>2</sub>, PPh<sub>3</sub>, TBHP, PhCl (pioneer work)
- FeCp<sub>2</sub>, TBHP, CH<sub>3</sub>CN, 8 examples, 30-78% yield
- [RuCl<sub>2</sub>(*p*-cymene)<sub>2</sub>]<sub>2</sub>, CH<sub>3</sub>CN, TBHP, 12 examples, 27-78% yield
- Cu(0), Selectfluor, CH<sub>3</sub>CN, 25 examples, 30-95% yield

Non-metal CDC

- TBAB, H<sub>2</sub>O, TBHP, air, 20 examples, 57-89% yield
- CBr<sub>4</sub>, O<sub>2</sub>, neat, 20 examples, 35-82% yield

Scheme 11. Methodologies for the synthesis of xanthenes through intramolecular cross-dehydrogenative coupling (CDC) of 2-aryloxybenzaldehydes

a cyclic seven membered transition state (**29**) by addition of TBHP followed by reductive elimination to afford the desired compounds **30** <sup>161</sup>.

Over the last years, the development of green CDC methodologies has sprout in order to compete with these transition-metal catalysed approaches (Scheme 11) and to address some drawbacks towards their use in the pharmaceutical industry (metal harsh elimination and consequent toxicity). Rao's et al. <sup>164</sup> reported a metal-free tetrabutylammonium bromide (TBAB)-promoted intramolecular annulation in aqueous medium, via oxidative coupling of C–H bond/aromatic C–H of several 2-aryloxybenzaldehydes. After reaction conditions optimization, this reaction tolerated both electron-donating and electron-withdrawing functional groups (Scheme 11). Interestingly, when 2-(4'-phenoxyphenoxy)-benzaldehyde was employed, the formyl C–H bond selectively coupled with the 2-phenoxy part to give the phenoxy-9*H*-xanthene-9-one while leaving the 4'-phenoxy part intact, attesting that the intramolecular route is favored by the oxidative coupling protocol.


In a related investigation, Tang et al. <sup>165</sup> developed a straightforward ring closure protocol for the synthesis of xanthenes from 2-aryloxybenzaldehydes through a simple and practical carbon tetrabromide promoted intramolecular aerobic oxidative dehydrogenative coupling reaction (Scheme 11). The scope and versatility of this reaction was also investigated, by experimenting different solvents, halogen reagents, and substituents, with good tolerance for different functional groups, being inclusively extended for the synthesis of fluorenones by using 2-arylbenzaldehydes as substrates. After preliminary studies on the reaction mechanism, the authors also suggested that the reaction may proceed through a radical pathway <sup>165</sup>.

**4.2.2 Cyclization of diaryl ethers to xanthenes by an acyl radical cyclization.** Although CDC is the most frequently used methodology for the synthesis of xanthenes from diaryl ethers, other less explored methodologies, with substrates containing a carboxylic acid derivative moiety, have recently been reported. An efficient intramolecular radical cyclization reaction via photoredox catalysis was developed for the synthesis of dibenzocycloketone derivatives using methylene blue as a


photosensitizer (Scheme 12A) <sup>166</sup>. This strategy relies on the fact that carboxylic acids are attractive and widely used synthons to prepare acyl radicals by decarboxylation through photoredox catalysis. The authors used 2-(2-phenoxyacetyl) benzoic acid as a model substrate to obtain the corresponding eight-membered dibenzocycloketones and, after reaction conditions optimization, the scope of the reaction was extended to other six, seven and eight-membered dibenzocycloketones. Xanthone (**63**) was obtained in 75% yield through reaction of 2-phenoxybenzoic acid (**62**) with methylene blue, PPh<sub>3</sub> and 2,4,6-collidine in 2 mL DMA and irradiated with LED white light at 25 °C for 24 h <sup>166</sup>.

#### 4.2.3 Cyclization of diaryl ethers to xanthenes by a Friedel–Crafts acylation. A promising metal-free variant of the intramolecular


##### A. Cyclization of diaryl ethers to xanthenes by an acyl radical cyclization




##### B. Cyclization of diaryl ethers to xanthenes by a Friedel–Crafts acylation



##### C. Cyclization of diaryl ethers to xanthenes by a FeCl<sub>3</sub> mediated direct intramolecular acylation of esters



##### D. Cyclization of diaryl ethers to xanthenes by palladium-catalyzed carbonylation/C–H activation



a) Y = N<sub>2</sub><sup>+</sup> BF<sub>4</sub><sup>−</sup>

R<sup>1</sup> = H, CH<sub>3</sub> or OCH<sub>3</sub>; R<sup>2</sup> = H or CH<sub>3</sub>; R<sup>3</sup> = H, CH<sub>3</sub>, OCH<sub>3</sub>, *t*Bu or Cl  
 Pd(PPh<sub>3</sub>)<sub>4</sub>, K<sub>2</sub>CO<sub>3</sub>, TBAB, CO, 80 °C  
 (23 examples, 25-90% yield)

b) Y = I

R<sup>1</sup> = H or CH<sub>3</sub>; R<sup>2</sup> = H; R<sup>3</sup> = H, CH<sub>3</sub>, OCH<sub>3</sub>, *t*Bu, Cl, F or CHO  
 Pd(OAc)<sub>2</sub>, P(Cy)<sub>3</sub>, PivONa.H<sub>2</sub>O, PivOH, DMSO, CO, 0.4 MPa, 90 °C, 5 h  
 (8 examples, 35-71% yield)

Scheme 12. Methodologies used for the cyclization of diaryl ethers to xanthenes

Friedel–Crafts acylation reaction, that employs mild conditions and avoid excesses of harsh acids, was described by Motiwala et al. <sup>167</sup>. This simple strategy consists in dissolving readily available acid chlorides in 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP), a strong hydrogen-bond-donating solvent, and allows the reaction to stir at room temperature for 4 h (Scheme 12B). The reaction was optimized using 4-(3,4-dimethoxyphenyl)butanoic acid as a model against solvent, HFIP loading, and reaction time to obtain 6,7-dimethoxy-1-tetralone. After optimal conditions were obtained, the scope was extended to a range of structurally diverse carboxylic acids; the



## REVIEW

## Organic Chemistry Frontiers

formation of six-membered rings was generally favored over five- and seven-membered rings being xanthone **65** obtained in 77% yield and 1,2,3-trimethoxy-9*H*-xanthen-9-one (**67**) in 93% yield (Scheme 12B).

**4.2.4 Cyclization of diaryl ethers to xanthenes by a FeCl<sub>3</sub> mediated direct intramolecular acylation of esters.** Benzyl esters have also shown to be suitable building blocks for the synthesis of xanthone derivatives. This serendipitous example of cyclization of diaryl ethers bearing a carboxylic acid moiety emerged as a way to dodge the disadvantages sometimes associated with the classical methods<sup>168</sup>. After an initial screening concerning several catalysts, dichloromethyl methyl ether loading, and reaction times, and with the optimal conditions in hands, the scope of the reaction was investigated and it was observed that the acylation could tolerate various functional groups such as OCH<sub>3</sub>, NO<sub>2</sub>, CO<sub>2</sub>CH<sub>3</sub>, Br, and Cl, affording the desired xanthenes in good yields (Scheme 12C). Mechanistic studies indicated that the formation of an intermediate, which results from the cooperatively activation of benzyl esters by FeCl<sub>3</sub> and Cl<sub>2</sub>CHOCH<sub>3</sub>, plays a key role in the acylation of esters.


**4.2.5 Cyclization of diaryl ethers to xanthenes by palladium-catalysed carbonylation/C–H activation.** Although the previous examples approached the use of diaryl ethers bearing a carboxylic acid moiety, this is not mandatory and the carbonyl bridge between the two aryl rings can be formed with the aid of carbon monoxide. Thus, palladium catalysed carbonylations/C–H activation reactions

emerged as an alternative methodology for the synthesis of xanthenes in the last years. The employment of a palladium catalyst (and ligand), base/additive, and carbon monoxide proceeds smoothly when using 2-iodo diaryl ethers or the corresponding diazonium salts to form the desired xanthenes, with great tolerance for different functional groups (Scheme 12D). The proposed mechanism of the intramolecular cyclization is similar in both strategies and proceeds through the initial oxidative addition of Pd, insertion of CO, activation of an *ortho* aromatic C–H with the assistance of the base, subsequent reductive elimination to afford the intended xanthone, and regeneration of Pd for the next catalytic cycle.


### 4.3 Synthesis of azaxanthenes

The synthesis of azaxanthenes via the diaryl ether route requires the synthesis of an intermediate, typically a cyanopyridyl phenyl ether (Scheme 13) or a phenoxy nicotinic acid derivative (Scheme 14). Cyanopyridyl phenyl ether derivatives can be obtained either through S<sub>N</sub>Ar of aromatic halides with hydroxypyridine or phenol derivatives. Phenoxy nicotinic acids are also achieved via S<sub>N</sub>Ar, from building blocks containing the carboxylic acid moiety or other functionalities (e.g. methyl) that are easily converted into the desired carboxylic acid. Other strategies include the use of quinolinimide as starting material, which can be further transformed into the corresponding carboxylic acid (Scheme 14). An excess of protic acid (often as solvent) such as sulfuric


#### A. Synthesis of 2-azaxanthone from 4-chloro-3-nitropyridine and phenol



#### B. Synthesis of 1-azaxanthone from 3-pyridinol and bromobenzene



#### C. Synthesis of 7-bromo-2-chloroazaxanthone from 3-chloropyridine and 4-bromophenol



Scheme 13. Synthesis of azaxanthenes via cyanopyridyl phenyl ether derivatives (diaryl ether route)

## REVIEW

## Organic Chemistry Frontiers

acid or polyphosphoric acid (PPA) is usually the reagent of choice for the intramolecular Friedel-Crafts cyclization reaction.

#### 4.3.1 Synthesis of azaxanthenes via cyanopyridyl phenyl ether.

The synthesis of azaxanthenes via cyanopyridyl phenyl ether is a very straightforward methodology, since the cyclization step is easily achieved due to the presence of a pendent cyano moiety. 2-Azaxanthone (**44**) was synthesized via this methodology by Kruger et al.<sup>169</sup>, starting from the condensation of 4-chloro-3-nitropyridine (**72**) with phenol to yield intermediary 3-nitro-pyridyl phenyl ether (**73**) (Scheme 13A). Reduction of the nitro group, oxidative conversion of the amine to the corresponding nitrile, and further cyclization with sulfuric acid at 195 °C led to 2-azaxanthone (**44**), in very low yield (1% over 4 steps), probably due to the formation of sulfonated phenoxy derivatives that were soluble in water. A similar strategy was used for the synthesis of 1-azaxanthone (**80**)<sup>78</sup> (Scheme 13B). 3-Phenoxy-pyridine (**77**) was used as starting material, previously obtained from the reaction between 3-pyridinol (**76**) and bromobenzene. *N*-oxide formation with *m*-chloroperbenzoic acid, and subsequent reaction with dimethyl sulfate produced *N*-methoxymethyl sulfate salts. Due to the fact that the resulting salts are generally hygroscopic and difficult to handle, reaction with sodium cyanide under nitrogen atmosphere, yielded stable 2-cyano-3-phenoxy-pyridine (**79**). 1-Azaxanthone (**80**) was obtained via direct ring closure in 17% overall yield in four steps from 3-pyridinol (**76**). A di-halogen-substituted azaxanthone was also synthesized via cyanopyridyl phenyl ether from 3-chloropiconitrile (**81**) and 4-bromophenol (**82**) (Scheme 13C). The diaryl ether **83** was obtained through *S<sub>N</sub>Ar* and further intramolecular Friedel-Crafts reaction in PPA and an *in situ* hydrolysis yielded the 1-azaxanthone core **84**. Nitrogen oxidation and treatment with phosphoryl chloride originated the desired chloride **85** with 62% overall yield, considering 4 reaction steps (Scheme 13)<sup>149</sup>.


**4.3.2 Synthesis of azaxanthenes via phenoxynicotinic acid derivatives.** 4-Azaxanthone (**43**), also known as 9-oxa-4-azaxanthone or 5*H*-[1]benzopyrano[2,3-*b*]pyridin-5-one, was the first azaxanthone to ever be synthesized, following a methodology involving phenoxynicotinic acid derivatives

(Scheme 14A). Preparation of 2-aminonicotinic acid (**87**) via Hofmann reaction on quinolinimide **86**, previously obtained from 8-hydroxyquinoline and further treatment of **87** with nitrous oxide yielded 2-hydroxynicotinic acid (**88**). Transformation into 2-chloronicotinic acid (**30**) and condensation of the corresponding sodium salt with sodium phenoxide yielded 2-phenoxynicotinic acid (**89**) which, after cyclization afforded compound **43**<sup>170</sup>. Another method for the synthesis of azaxanthenes was developed by Villani and collaborators<sup>78</sup>, taking phenoxy-pyridinecarboxylic acids as starting materials, with the nitrogen in different positions, according to the desired azaxanthone. The reactions employed allowed the synthesis of azaxanthenes previously described in better yields. For the synthesis of 4-azaxanthone (**43**, Scheme 14B), 2-chloronicotinic acid (**30**) reacted with phenol in the presence of sodium methoxide to yield 2-phenoxynicotinic acid (**89**). The ring closure was achieved with the use of phenylphosphonic acid (PPA), in 83% overall yield [2 steps from 2-chloronicotinic acid (**30**). For the synthesis of 3-azaxanthone (**93**, Scheme 14C), the starting materials used were 3-hydroxy-4-picoline (**44**) and bromobenzene, which through the Ullmann condensation led to 3-phenoxy-4-picoline (**90**). 3-Phenoxyisonicotinic acid (**92**) was obtained by oxidation of **91** and heating of **92** with PPA, similarly to the previous reaction, led to 3-azaxanthone (**93**) in 16% overall yield, in three steps from 3-hydroxy-4-picoline (**90**)<sup>78</sup>. On the other hand, 2-azaxanthone (**44**) was prepared by using 4-nitro-3-picoline 1-oxide (**94**) and phenol as starting materials. Displacement of the nitro group by phenoxide in the presence of K<sub>2</sub>CO<sub>3</sub> gave phenyl ether **95** that was *N*-deoxygenated to the pyridyl ether **96**. Oxidation gave the desired 4-phenoxynicotinic acid (**97**) which after ring closure gave 2-azaxanthone (**44**) in 21% overall yield in 4 steps from 4-nitro-3-picoline 1-oxide (**94**) and phenol (Scheme 14D)<sup>78</sup>. It is also worth to highlight a simple, clean methodology for the synthesis of azaxanthenes and azathioxanthenes. Li et al.<sup>171</sup> developed a synthetic route that did not use expensive materials and had solvent-free conditions, starting from phenoxy-pyridine acids. In order to infer the best catalyst, studies with a series of metal triflates were carried out. It was concluded that ytterbium(III) trifluoromethanesulfonate was the


REVIEW

Organic Chemistry Frontiers


A. Synthesis of 4-azaxanthone from quinolinimide




B. Synthesis of 4-azaxanthones from 2-chloronicotinic acid and phenol




C. Synthesis of azaxanthones from 3-hydroxy-4-picoline and bromobenzene



D. Synthesis of azaxanthones from 4-nitro-3-picoline 1-oxide and phenol



E. Synthesis of azaxanthones via phenoxynicotinic acid derivatives



Scheme 14. Synthesis of azaxanthones via phenoxynicotinic acid derivatives (diaryl ether route)

best catalyst, and that the use of a Brønsted acid would slightly improve the yield in the synthesis of 4-azaxanthone and derivatives (Scheme 14E).

**4.3.3 Synthesis of azaxanthones via amide derivatives.** In an effort to find new  $\beta$ -amyloid precursor protein cleaving enzyme 1 (BACE1) inhibitors, new azaxanthones were synthesized<sup>149</sup>. Compared to previously synthesized xanthones, the introduction of a nitrogen could improve in vitro potency, cardiovascular safety, pharmacokinetics, drug metabolism, and modulate central nervous system penetration<sup>172</sup>. These

features could also be evidenced by a fluorine-substitution at the 3- or 4-position. The combination of these features would ultimately lead to an overall improvement of the compounds as BACE1 inhibitors. Since these molecules had never been synthesized before, different synthetic approaches were attempted. For the synthesis of 3-aza-4-fluoroxanthone **104** (Scheme 15), the first step was the regioselective copper-catalysed Ullmann coupling between 2,5-dibromobenzoic acid (**100**) and 2-fluoro-hydroxypyridine (**101**), which originated a diaryl ether **102**. This intermediate **102** was treated with diethylamine and 2-(1H-benzotriazole-1-yl)-1,1,3,3-


tetramethylammonium tetrafluoro-borate (TBTU) to give amide **103**. An amide-directed lithiation on the pyridine, followed by an *in situ* cyclization furnished a fluorinated azaxanthone **104** [30% overall yield in 3 reaction steps from 2,5-dibromobenzoic acid (**100**)]. From the same amide intermediate **103**, *N*-oxidation was performed on the pyridine nitrogen (**105**) and subsequent treatment with phosphoryl chloride afforded a chloride derivative **106**, which produced a tri-halogenated azaxanthone **107** after a tandem lithiation/cyclization (Scheme 15), with an overall yield of 9% in five steps from 2,5-dibromobenzoic acid (**100**) and 2-fluoro-hydroxypyridine (**101**) (Scheme 15) <sup>149</sup>.

## 5. SYNTHESIS OF XANTHONES VIA CHROMEN-4-ONE DERIVATIVES


The chromen-4-one core constitutes a versatile building block that can be used for the synthesis of xanthone derivatives. To form the additional aromatic ring, several strategies can be applied, mainly the traditional Diels-Alder or other cycloaddition reactions. Depending on the substitution pattern of the chromen-4-one derivative, this section will be subdivided in three subsections: synthesis of xanthones from 2-substituted (section 6.1), 3-substituted (section 6.2) and 2,3-disubstituted chromen-4-ones (section 2.3) (Scheme 16).

### 5.1 Synthesis of xanthones from 2-substituted chromen-4-ones

**5.1.1 Via Diels-Alder reaction.** The high versatility and efficiency of Diels-Alder reactions have also been extended to the synthesis of xanthones from 2-substituted chromen-4-ones. Xanthone-1,2,3-triazole dyads were obtained by two different approaches, both starting from (*E*)-2-(4-arylbut-1-en-3-yn-1-yl)chromones, previously obtained by a base-catalysed aldol reaction of 2-methylchromone and arylpropargyl aldehydes <sup>173</sup>. While the first approach adopted a methodology in which the xanthone core is primarily constructed using chromones as dienes in the Diels-Alder reaction, followed by the construction of the triazole ring, in the second strategy the process was inverted, starting with the triazole ring followed by the construction of the xanthone core (Scheme 17A). Thus, in the first methodology, a Diels-Alder reaction of the chromone derivatives **108** with *N*-methylmaleimide under microwave irradiation gave Diels-Alder adducts **109**. Oxidation of the obtained adducts with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) gave xanthone derivatives **110** which, upon reacting with sodium azide in refluxing DMF provided the desired xanthone-1,2,3-triazole dyads **111** in excellent yields (88-90%). Second approach consisted in 1,3-cycloaddition reaction of chromones **108** with sodium azide to form the triazole derivatives (**112**) followed by methylation of the triazole NH group prior to Diels-Alder reaction with *N*-methylmaleimide and oxidation with DDQ to provide the xanthone-1,2,3-triazole



Scheme 15. Synthesis of azaxanthones via amide derivatives (diaryl ether route)



Scheme 16. Synthesis of xanthones via chromen-4-ones

## REVIEW


## Organic Chemistry Frontiers

dyads (**111**) in moderate to good yields (57-80%)<sup>173</sup>. (*E*)-2'-Propargyloxy-2-styrylchromones (**115**) were used as substrates in a microwave-assisted intramolecular Diels–Alder reactions, affording chromeno[3,4-*b*]-xanthenes **116** (Scheme 17B)<sup>174</sup>. During optimization processes, besides the variation of solvent, Lewis acid, temperature, and reaction times, different oxidizing agents were also tested in order to push the reaction towards the oxidized product **116**, without the formation of secondary products that are not fully oxidized. Optimal conditions for MAOS of chromeno[3,4-*b*]xanthenes are the use of 1,2,4-

trichlorobenzene (1,2,4-TCB) as a solvent, under MW irradiation at 200 °C for 60 min and then the addition of the oxidizing agent chloranil to the reaction crude, under MW irradiation at 80 °C for 30 min<sup>174</sup>.

**5.1.2 Via electrocyclization.** Other example of transforming 2-substituted chromen-4-ones into xanthone derivatives is through an electrocyclization and subsequent oxidation (Scheme 18). The substrates 2-[(1*E*,3*E*)-4-arylbuta-1,3-dien-1-yl]-4*H*-chromen-4-

### A. Synthesis of xanthone-1,2,3-triazole dyads




#### Experimental conditions:

a) *N*-methylmaleimide, focused MW, 10 min, 200 °C  
b) DDQ, toluene, 1h, 100 °C


c) NaN<sub>3</sub>, dry DMF, 1h, reflux  
d) K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>SO<sub>4</sub>, acetone, 1h, reflux  
e) *N*-methylmaleimide, focused MW, 10 min, 200 °C

### B. Synthesis of chromeno[3,4-*b*]-xanthenes through a MW-assisted intramolecular Diels–Alder reaction



Scheme 17. Synthesis of xanthenes from 2-substituted chromen-4-ones via Diels–Alder reaction

## REVIEW



Scheme 18. Synthesis of xanthenes through an electrocyclization and subsequent oxidation


ones **117** are previously obtained in good yields by condensation of 2-methyl-4*H*-chromen-4-ones with cinnamaldehydes in the presence of sodium ethoxide. Thus, the target 1-aryl-9*H*-xanthen-9-ones **118** are obtained under refluxing 1,2,4-TCB in the presence of a catalytic amount of iodine and subsequent demethylation with  $\text{BBr}_3$ , in moderate to good yields<sup>175</sup>. This study also disclosed that some of the tested compounds had improved scavenging activity when compared with previously reported analogues, being promising pharmacophores with potential therapeutic applications associated to oxidative stress disorders<sup>175</sup>.

## 5.2 Synthesis of xanthenes from 3-substituted chromen-4-ones

**5.2.1 Via a tandem [4+1]-[4+2] cycloaddition.** Bornadiego et al.<sup>176-178</sup> reported the synthesis of 4-aminoxanthenes via a tandem [4+1]-[4+2] cycloaddition of 3-carbonylchromones, isocyanides and dienophiles (Scheme 19A). The authors were able to extend the reaction scope to different dienophiles, either symmetric *N*-phenyl- and *N*-methylmaleimide, maleimide, and maleic anhydride<sup>176</sup>, or asymmetric ones like acrylonitrile or methyl vinyl ketone<sup>177</sup>. Additionally, they were also able to extend this methodology to carbonylchromones containing both electron-withdrawing and electron-donating groups<sup>178</sup>. Mechanistically, 4-aminoxanthone is obtained through a [4+1] cycloaddition of a 3-carbonylchromone **119** with an isocyanide **120** to give an iminolactone intermediate **121**. Tautomerization to aminofuran **122**, which suffers a [4+2] cycloaddition with the dienophile **123**, leads to 7-oxabicyclo [2.2.1]heptane (**124**). This is readily transformed to 1-hydroxydihydroxanthone **125** by the opening of the oxygen bridge assisted by the nitrogen lone pair. Dehydration finally affords the aromatic 4-aminoxanthone **126** (Scheme 19A)<sup>178</sup>.


**5.2.2 Via a tandem Heck reaction/double C-H activation/retro Diels-Alder pathway.** Another strategy towards the

### A. Synthesis of 4-aminoxanthenes via a tandem [4+1]-[4+2] cycloaddition




### B. Synthesis of xanthenes via a palladium-catalyzed cascade reaction of 3-iodochromones with aryl iodides

#### Scope of Aryl Iodides



#### Scope of Iodochromones



Scheme 19. Synthesis of xanthenes from 3-substituted chromen-4-ones



## REVIEW

## Organic Chemistry Frontiers

construction of the xanthone core from 3-substituted chromen-4-ones is *via* a tandem Heck reaction/double C-H activation/retro Diels-Alder pathway. The use of readily available 3-iodochromones **127** and **130**, aryl iodides **128** and **131**, and norbornadiene as starting materials in this palladium-catalysed cascade reaction enables access to diverse annulated xanthenes (**129** and **132**, Scheme 19B) <sup>179</sup>. This process is initiated by oxidative addition of Pd(0) to the chromone ring system, followed by norbornadiene insertion and C-H activation of the 2-position of the chromone ring. A second oxidative addition of the aryl iodide, subsequent reductive elimination, and second C-H activation formed the desired xanthenes <sup>179</sup>.

**5.2.3 Synthesis of azaxanthenes by reaction of carbonitrile chromones and oximes.** To better understand condensation reactions of 4-oxo-4*H*-[1]benzopyran-3-carbonitriles, Ghosh et al. <sup>180</sup> have performed reactions that yielded, among other compounds, azaxanthenes. The starting carbonitrile chromone **133** reacted with acetylacetone (**134**), in the presence of piperidine, furnishing a 4-azaxanthone derivative **135** in 80% yield. The mechanism through which this reaction occurs was investigated, and it was concluded that **134** undergoes Michael addition to the  $\alpha,\beta$ -unsaturated ketone. Concomitantly, the pyrone ring opens, yielding an intermediate which undergoes, probably in a concerted manner, a double cyclisation (Scheme 20A). The same study also contemplated the synthesis of the same compound (**135**), taking an oxime (**137**) as starting material (Scheme 20A). Given that the chromone-3-nitriles are prepared from oximes, their use for preparing azaxanthenes shortens the synthetic route, although the yield, 67%, was not as good as when the nitrile was used as starting material <sup>180</sup>.

**5.2.4 Synthesis of azaxanthenes by reaction of a cyano chromone and malononitrile.** Reaction of cyano chromone **133** with malononitrile led to **138** which was then deaminated by sodium nitrite in trifluoroacetic acid. Further chlorination using phosphoryl chloride and phosphorus pentachloride afforded **139** in 38% yield, which was converted to compound **140** by catalytic hydrogenation (Scheme 20B), in 28% overall yield from cyano chromone **133** in four reaction steps <sup>181</sup>.

**5.2.5 Synthesis of azaxanthenes via a one-pot condensation between 1,3-bis-silyl enol ethers with 3-cyanobenzopyrylium triflates.** Azaxanthone derivatives could also be attained by domino reactions in a one-pot process. The approach used by Langer and Appel <sup>182</sup> consisted on the condensation of 1,3-bis-silyl enol ethers (**143**), unsymmetrical acetoacetone, with 3-cyanobenzopyrylium triflates (**142**). This yielded an open-chained product (**143**), which were treated with triethylamine, to give an azaxanthone (**144**) in 46% overall yield, through a domino “retro-Michael-lactonization-aldol” reaction (Scheme 20C).


**5.2.6 Synthesis of azaxanthenes by reaction of chromone-3-carbaldehyde with  $\alpha$ -aminoacids.** In an attempt to synthesize (hydroxybenzoyl)pyrroles, Figueiredo et al. <sup>183</sup> have discovered an azaxanthone, from a chromone moiety. The reaction started with a chromone aldehyde **145** and ethyl glycinate hydrochloride, and depending on the reaction conditions, several compounds were formed. In order for the azaxanthone derivative **146** to be synthesized, in 12% overall yield, one equivalent of methyl glycinate and 0.5 equivalent of potassium carbonate were required, in refluxing toluene. The authors proposed a mechanism for the formation of the azaxanthone intermediate, which comes from the oxidation of the dihydroxanthone (Scheme 20D).

**5.2.7 Synthesis of azaxanthenes via a one-pot Michael addition between an alkynylchromone and an aldimine.** One-pot approaches were also an efficient method of obtaining azaxanthenes. In this scope, the starting material chosen was 2-(1-alkynyl)-2-alken-1-ones (Scheme 20E), which are interesting for multicomponent reactions, as they are prone to transition metal, Lewis acid, and electrophile-induced cascade cyclization reactions promoted by nucleophiles. Since these starting compounds have previously been reported to be involved in reactions initiated by Michael addition of nitrogen, it was postulated that *N*-unsubstituted aryl aldimines would act as nucleophiles in Michael addition reactions with 3-(1-alkynyl)chromones. Subsequently, a cascade sequence would be initiated, ultimately leading to the production of azaxanthenes <sup>14</sup>. The optimal conditions for this reaction were also investigated. DMF was found to be the best solvent, against ethanol, DMSO, toluene, and dioxane. As for the base, it was shown that three equivalent of ammonium acetate provided the optimal conditions, but the reaction also occurred with ammonium carbamate, ammonium bicarbonate, and ammonium formate. In the opposite way, ammonium chloride and ammonium sulfate did not yield the desirable product <sup>14</sup>. The same study also described the synthesis of 2,4-diazaxanthenes (**151**) through the same methodology (Scheme 20F). In an effort to extend the cascade reaction to other Michael acceptors, 3-cyanochromone (**141**) was chosen as starting material. Testing different reaction conditions, it was demonstrated that the use of copper(II) chloride as an oxidant would lead to yields up to 85%. For the production of the unsubstituted 2,4-diazaxanthone (**151**), the aldehyde employed was paraformaldehyde (**150**), and the yield was 74% (Scheme 20F) <sup>14</sup>.


**5.2.8 Synthesis of azaxanthenes via a domino reaction using indium as a catalyst.** A green approach for the formation of azaxanthenes was proposed by Poomathi et al. <sup>184</sup>. This group managed to synthesize azaxanthenes through a domino reaction using a small amount of catalyst and ethanol as solvent. Studies on the catalyst showed that indium triflate was the best choice of a Lewis acid for this reaction, and that it could be recovered after the reaction and maintain its catalytic efficiency for, at least, five reactions.

## REVIEW


## A. Synthesis of azaxanthenes by reaction of carbonitrile chromones and oximes




## B. Synthesis of azaxanthenes by reaction of a cyano chromone and malononitrile



## C. Synthesis of azaxanthenes via a one-pot condensation between 1,3-bis-silyl enol ethers with 3-cyanobenzopyrylium triflates



## D Synthesis of azaxanthenes by reaction of chromone-3-carbaldehyde with α-aminoacids




Scheme 20. Synthesis of xanthenes from 3-substituted chromen-4-ones


## REVIEW

## Organic Chemistry Frontiers


### E. Synthesis of azaxanthenes via a one-pot Michael addition between an alkynylchromone and an aldimine



### F. Synthesis of 2,4-diazaxanthone 127



### G. Synthesis of azaxanthenes via a domino reaction using indium as a catalyst



Scheme 20 (cont.). Synthesis of xanthenes from 3-substituted chromen-4-ones

In order to obtain azaxanthenes (**154**), 3-formylchromone (**152**) reacted with *N,N'*-dibutyl-2-nitroethene-1,1-diamine (**153**), using In(OTf)<sub>3</sub> in ethanol, in reflux, as can be seen in Scheme 20G. This reaction allowed the synthesis of a series of derivatives, whose yields ranged from 80% to 95%<sup>184</sup>. The reaction mechanism hypothesized involves a Henry reaction between 3-formylchromone (**152**) and *N,N'*-dibutyl-2-nitroethene-1,1-diamine (**153**) (Scheme 20G). Elimination of water and further 6π-electrocyclization, yields azaxanthenes **154**<sup>184</sup>.

### 5.3 Synthesis of xanthenes from 2,3-disubstituted chromen-4-ones

**5.3.1 Via photo-induced tandem cyclization 3-iodoflavones with five membered heteroarenes.** An excellent example that provides rapid access to polycyclic xanthenes without the requirement of any transition-metal-catalyst and/or oxidant additives is a tandem cyclization via the coupling of 3-iodoflavones with five-membered heteroarenes<sup>185</sup>. This procedure allowed the preparation of a broad variety of novel polycyclic xanthone frameworks with excellent regioselectivity in good yields under mild and environmentally friendly reaction conditions. Mechanistic studies unveiled that the reaction proceeds *via* two consecutive C–C bond formations after irradiation of the substrate and *N*-methylpyrrole in CH<sub>3</sub>CN using a high-pressure mercury lamp (500 W) at room temperature for 7 h (Scheme 20A). It was possible to observe


that obtained yields were lower when in the presence of electron-donating groups on both R<sup>1</sup> and R<sup>2</sup> and higher when in the presence of only one electron-donating group at either R<sup>1</sup> or R<sup>2</sup>. The substrate scope was also extended to other five-membered heteroarenes, with the desired xanthone derivatives being obtained in lower yields<sup>185</sup>.

**5.3.2 Via intramolecular [4+2] cyclization.** Another procedure that allows the preparation of xanthone derivatives without the need for a transition metal catalyst, consists in a phase transfer reagent promoted tandem ring-opening and ring-closing reaction of 3-(hepta-1,6-diyn-1-yl)chromone derivatives followed by a intramolecular [4+2] cyclization<sup>186</sup>. The use of *t*-BuOK, *n*-Bu<sub>4</sub>NCl and phenylacetonitrile as an anion transfer reagent in DMSO at 110 °C for 20 min under MW irradiation proceeded smoothly for the preparation of several xanthone derivatives (Scheme 21B), with the exception of R<sup>2</sup> = H, and R<sup>1</sup> = Ph, being the latter probably due to steric effects of two phenyl groups at the *ortho*-position<sup>186</sup>.


**5.3.3 Via Friedel-Crafts alkylation.** An intramolecular Friedel Crafts alkylation strategy using AlCl<sub>3</sub> in dichloromethane was used to smoothly synthesize the corresponding benzoxanthenes in low to moderate yields (Scheme 21C), with the more electron rich aromatic systems in the 2-position of the chromone being obtained with increased efficiency<sup>187</sup>.

## REVIEW


## A. Synthesis of xanthenes via photo-induced tandem cyclization 3-iodoflavones with five membered heteroarenes



## B. Synthesis of xanthenes via intramolecular [4+2] cyclization



## C. Synthesis of xanthenes via Friedel-Crafts alkylation




Scheme 21. Synthesis of xanthenes from 2,3-disubstituted chromen-4-ones

**5.3.4 Synthesis of azaxanthenes by reaction of 4-oxo-4H-1-benzopyran-3-carbonitriles and aminoacrylate or ethyl propiolate.** While performing studies on potential anti-anaphylactic agents, Nohara *et al.* applied a series of different methods to synthesize azaxanthone derivatives, inspired by xanthone-2-carboxylic acids bearing a tetrazole group, which have proven oral antiallergic activity<sup>181, 188</sup>. The authors converted 4-oxo-4H-1-benzopyran-3-carbonitriles **161** into 2-amino-4-oxo-4H-1-benzopyran-3-carboxaldehydes **162** which were found to be good starting materials for the synthesis of some types of heterocycles. After the synthesis of these intermediates, some different synthetic methodologies were applied to produce different azaxanthenes (Scheme 22). The first method used was the reaction of **162** with ethyl propiolate with triethylamine (TEA) in DMF, yielding an aminoacrylate **163** (Scheme 22A), which was then converted by further heating to the desired azaxanthone **164**, in 19% yield. A

similar product could be obtained by a one-step reaction, using methyl malonyl chloride (Scheme 22B), in 36% yield<sup>181</sup>. A different approach was the reaction of **162** with cyanoacetylene, yielding 3-cyano derivatives (**167**), in 36% yield (Scheme 22D). In this reaction, no basic catalyst was needed. The search for an alternative to this methodology, since cyanoacetylene is unstable, sublimates at low temperatures, among other undesirable properties, led to the synthesis of the same type of compounds **167** in 33% yield, using  $\alpha$ -chloroacrylonitrile with triethylamine (Scheme 22E). A second alternative to prepare **167** is the treatment of **162** with DMF and cyanoacetyl chloride, which produced the desired compound in 49% yield (Scheme 22F)<sup>181</sup>. The change of DMF for dichloromethane, in the reaction with cyanoacetyl chloride, originated an amide intermediate (**168**) (Scheme 22G), which could be converted into an azaxanthone **169** by heating with pyridine.

REVIEW

Organic Chemistry Frontiers



Scheme 22. Syntheses of azaxanthenes starting from carbalddehyde 66

In this case, the azaxanthone obtained was 2-hydroxy substituted (**169**), in 22% yield<sup>181</sup>. The same group used another synthetic strategy towards the synthesis of azaxanthenes, which was the condensation of **162** with malonaldehyde. As such, a derivative of **162** was chosen to react with malonaldehyde bis(dimethyl acetal), in the presence of formic acid in boron trifluoride etherate, originating compound **165** in 61% yield (Scheme 22C). Afterwards, an oxime (**166**) was prepared through the reaction of **165** with hydroxylamine, with 95% yield. Finally, a nitrile (**168**) was obtained by the reaction with phosphoryl chloride in DMF (Scheme 22), yielding 36% of the desired compound<sup>181</sup>. The same group also performed reactions where the final product would be a 2-substituted azaxanthone. The methods used ethyl cyanoacetate and piperidine (Scheme 22H, 95% yield), cyanoacetic acid and pyridine (Scheme 22I, 55%) and diethyl acetylenedicarboxylate (Scheme 22J, 91%), yielding azaxanthone derivatives **170**, **171** and **172**, respectively<sup>181</sup>.


### 5.3.5 Synthesis of azaxanthenes via a modified Friedländer reaction.

The synthesis of azaxanthenes can also be achieved


through modification of the Friedländer reaction. Using *o*-aminoaromatic aldehydes and unmodified ketones, Dormer *et al.* (2003) proceeded to synthesize 2-alkylsubstituted products, in a regioselective manner. The main objective was to evaluate the efficiency of several catalysts in terms of regioselectivity, conversion ratio and reaction conditions, and it was concluded that 1,3,3-trimethyl-6-azabicyclo[3.2.1]octane (TABO) was the most effective. Slow addition of the ketone had a positive impact on regioselectivity, and reaction temperature also played a key role<sup>189</sup>. For the preparation of an azaxanthone using this method, 2-amino-3-formylchromone (**162**) and 2-hexanone (**173**) were chosen as starting materials. The chromone was slowly added to a solution of the ketone, sulfuric acid, and the catalyst in ethanol. Out of all the reactions this group performed, this was the one with the lowest regioselectivity [80% 3-substituted (**174**) (79% overall): 20% 2,3-disubstituted (**175**) (20% overall)] Scheme 23A). Curiously, this reaction did not occur under the conventional Friedländer conditions<sup>189</sup>.

## REVIEW


## A. Synthesis of azaxanthenes via a modified Friedländer reaction



## B. Synthesis of azaxanthenes via a gold-catalyzed reaction



## C. Synthesis of azaxanthenes by reaction of the chromone-carbaldehyde with phosphonyl-containing alkynes



Scheme 23. Synthesis of azaxanthenes from 2,3-disubstituted chromen-4-ones

**5.3.6 Synthesis of azaxanthenes via a gold-catalysed reaction.** A collection of *N*-heterocycles, including azaxanthenes, was synthesized through gold catalysed synthesis. The condensation of 2-amino-3-formylchromone and acetophenone derivatives, using gold(III) chloride and silver hexafluoroantimonate(V), and a mixture of acetonitrile and methanol as solvents, gave aryl substituted azaxanthenes with good yield (Scheme 23B) <sup>190</sup>.

**5.3.7 Synthesis of azaxanthenes by reaction of the chromone-carbaldehyde with phosphonyl-containing alkynes.** The synthesis of the azaxanthone scaffold was also described by a different route in an attempt to prepare 2-difluoromethyl-azaxanth-3-ylphosphonates. Duda *et al.* <sup>191</sup> showed a synthetic method that led to these compounds through 2-amino-3-formylchromone (**172**) which reacted with unsymmetrical substituted electron-poor halogen- $\text{CF}_2$  and phosphonyl-containing alkynes **179** and **181**. The reaction occurred with

DMSO as a solvent and a basic catalyst, at room temperature. It was proven that the best conditions for the synthesis of fluorine, chlorine, difluoromethyl and trifluoromethyl derivatives **179** was DMSO as solvent and *i*PrNEt as catalyst, while for bromine derivatives **181**, the best basic catalyst was potassium carbonate and DMF was the best solvent (Scheme 23C). In formulated hypothesis of reaction mechanism, the base-activated 2-amino-3-formylchromones reacts in a regioselective manner through a Michael-type addition with the carbon containing the halogen- $\text{CF}_2$ , which is the most electrophilic. Formation of a dipolar allene, and subsequent intramolecular cyclization into the dihydropyridine derivative, leads, after dehydration, to azaxanthone. The polar solvent has upmost importance in this reaction, serving as a stabilizer and possibly interacting with the allene intermediate. An interpretation of the reaction mechanism is depicted in Scheme 23C <sup>191</sup>.



## 6. OTHER METHODOLOGIES

Other important and conceptually different approaches, which do not belong to any of the categories described above, were also reported, mainly focusing in tandem and/or rearrangement reactions. Although some of them start from readily available substrates<sup>192</sup>, other require complex starting materials that usually need to be prepared prior to the reaction<sup>193,194</sup>, but also capable of reaching unusual substituents/patterns of substitution in the xanthonic core.

### 6.1 Synthesis of xanthenes *via* decarbonylative coupling.

Salicylaldehydes can also be used as substrates in rhodium-catalysed reductive decarbonylative couplings to give symmetrically substituted xanthenes **184** (Scheme 24A), containing formyl, bromo, chloro, fluoro, methyl, acetyl, nitro, hydroxyl, and trimethoxy groups (10 examples)<sup>141</sup>. Interestingly, this reaction revealed to be site-selective since when 4-hydroxyisophthalaldehyde was used as a substrate, the corresponding 9-oxo-9H-xanthene-2,7-dicarbaldehyde was obtained, indicating preferential reaction of the formyl group *ortho* to hydroxyl during decarbonylative homo-coupling process<sup>141</sup>.

### 6.2 Synthesis of xanthenes *via* a tandem etherification-acylation of diaryliodonium salts with salicylates

Salicylates can also be employed for the synthesis of substituted xanthenes **187**, when combined with diaryliodonium salts<sup>195</sup>. An intermolecular etherification-acylation in the presence of Cu(OTf)<sub>2</sub> as catalyst in dichloroethane (DCE) produced the desired xanthenes in good yields (Scheme 24B). Mechanistically, the reactions are initiated by the etherification of diaryliodonium salts with salicylates, and followed by an intramolecular acylation.

### 6.3 Synthesis of xanthenes *via* an intramolecular Diels-Alder reaction involving 2-(1,2-dichlorovinyl) aryl dienones

An intramolecular Diels-Alder reaction where both of the pyrone and the second aromatic rings are forged in a single step was reported for the synthesis of xanthenes<sup>192</sup>. The scope of the route to several substituted xanthenes was tested by using different 2-hydroxyacetophenones (**189**) to prepare the corresponding 1,2-dichlorovinyl acetophenones (**190** and **192**) that react *in situ* with cinnamaldehyde (Scheme 24C). Similarly, the scope of the reaction was also examined using a variety of commercially available  $\alpha,\beta$ -unsaturated aldehydes (Scheme 24C) with the expected xanthenes **191** and **194** being obtained in modest yields (23 examples, 18–62%)<sup>192</sup>.

### 6.4 Synthesis of xanthenes *via* a tandem reaction of 2-(4-(2-haloaryl)-4-hydroxybut-1-ynyl)benzaldehydes

12H-Benzo[*b*]xanthene-12-ones **196** were synthesized using Jones reagent and K<sub>2</sub>CO<sub>3</sub> promoted tandem reaction of 2-(4-(2-haloaryl)-4-hydroxybut-1-ynyl)benzaldehydes **195** (Scheme

24D)<sup>193</sup>. The scope and limitation of this reaction were studied and it was found that in addition to 2-bromo substituted substrate, 2-fluoro or 2-chloro derivatives could also be obtained. Due to fluoride's great inductive effect, which accelerates the rate-determining step of the nucleophilic aromatic substitution step, it was observed that fluoro substituted substrate reacted faster than bromo or chloro. However, bromo substituted substrate was more reactive than 2-chloro since, as is well known, Br is a better leaving group than Cl. It was proposed that the overall transformation presumably proceeds through an initial tandem reaction of **195** with the formation of a 3-acyl-2-naphthol key intermediate, which should be simultaneously transformed into the corresponding phenoxide in the presence of K<sub>2</sub>CO<sub>3</sub>. Further intramolecular *O*-arylation occurs to give the expected 12H-benzo[*b*]xanthene-12-ones **196**.

### 6.5 Synthesis of xanthenes *via* Au(I)-catalysed Michael addition/6-*endo*-trig cyclization/aromatization cascade annulation


Another strategy used for the synthesis of 12H-benzo[*b*]xanthene-12-ones **198** is through a Michael addition/6-*endo* cascade cyclization/aromatization, extensible to the synthesis of benzoacridones (Scheme 24E)<sup>194</sup>. While the presence of electron-withdrawing groups on the methoxyvinyl phenyl ring led to production of the target xanthenes in higher yields, when they were present in the phenol ring poor yields were obtained, regardless of the substitutions on the methoxyvinyl phenyl ring<sup>194</sup>. The observed outcome of the reaction could be explained by the formation of a complex with the gold catalyst, activating the 1,3-diphenylprop-2-yn-1-one which is then attacked by the hydroxyl group in a 6-*endo*-dig manner. Photodeauration of the formed intermediate and further attack of the newly formed enol ether to the electron-rich methoxyl vinyl ether gave the final xanthone after isomerization and aromatization through the release of methanol.

### 6.6 Synthesis of xanthenes *via* a base-promoted insertion reaction


In a completely different approach, xanthenes and azaxanthenes were obtained through a base-promoted insertion and transition-metal-free reaction of isolated internal alkynes with  $\beta$ -diketone compounds (Scheme 24F)<sup>196</sup>. Studies on the scope of the reaction revealed that different leaving groups on the acetylenic ketone **199** were well tolerated, although 1-(2-bromophenyl)-3-phenylprop-2-yn-1-one (LG = Br) produced the target xanthone **201** in highest yield and greatest chemoselectivity. The corresponding acetyl aryl derivative **202** was obtained as secondary product (10% yield) of the reaction, which increased when an electron-withdrawing substituent (R<sup>2</sup> = 7-F) was employed.

## REVIEW

## A. Synthesis of xanthenes via decarbonylative coupling




## B. Synthesis of xanthenes via a tandem etherification-acylation of diaryliodonium salts with salicylates




## C. Synthesis of xanthenes via an intramolecular Diels-Alder reaction involving 2-(1,2-dichlorovinyl) aryl dienones


## Scope of acetophenones




## Scope of cinnamaldehydes



## D. Synthesis of xanthenes via a tandem reaction of 2-(4-(2-haloaryl)-4-hydroxybut-1-ynyl)benzaldehydes



## E. Synthesis of xanthenes via Au(I)-catalyzed Michael addition/6-endo-trig cyclization/aromatization cascade annulation




Scheme 24. Other methodologies for the synthesis of xanthenes


## REVIEW

## Organic Chemistry Frontiers

### F. Synthesis of xanthenes via a base-promoted insertion reaction



### G. Synthesis of xanthenes via sequential fluoride ion-promoted Fries-Type rearrangement and nucleophilic aromatic substitution



Scheme 24 (cont.). Other methodologies for the synthesis of xanthenes

### 6.7 Synthesis of xanthenes via sequential fluoride ion-promoted Fries-Type rearrangement and nucleophilic aromatic substitution

A mild and efficient reaction promoted by fluoride ion was used to prepare 1,8-disubstituted xanthenes **204** from phenyl benzoate derivatives **203** (Scheme 24G)<sup>197</sup>. The reaction proceeded via a Fries-type rearrangement to generate a benzophenone intermediate that immediately cyclizes into the expected xanthenes due to the presence of phenoxide and fluorine moieties at C-2 and C-2' positions. This methodology is particularly useful for the synthesis of xanthenes with steric hindrance around C-9, which are difficult to obtain through other methodologies.

### 6.8 Synthesis of azaxanthone-derivatives: the case of five-membered ring azaxanthenes

Different types of azaxanthenes have been synthesized. Noteworthy for their biological activities are the five-membered ring azaxanthenes. The incorporation of a pyrazole and a triazole moiety in the xanthone core can be achieved by two different methods. In the first method (Scheme 25A), an intramolecular  $S_NAr$  would be responsible for the construction of the central ring; in the second method (Scheme 25B), an anionic intramolecular ring closure takes place. Concerning the first method, 1-(benzyloxy)pyrazole (**205**) and 1-benzyloxy-1,2,3-triazole (**206**) were metalated, using butyllithium in THF, at -78 °C. The second step was slightly different for the two lithiated compounds: for the pyrazole derivative, transmetalation with zinc(II) iodide or chloride was performed, followed by cross-coupling 2-fluorobenzoyl chloride and tetrakis(triphenylphosphine)palladium(0) in THF at 20 °C. The next step was debenzoylation, using hydrochloric acid<sup>198</sup>. On the other hand, for the triazole derivative, the transmetalation step occurred with tributylstannane, and the debenzoylation with  $H_2/Pd$ <sup>199</sup>. The ring closure, which proceeded under mild conditions, using potassium carbonate in DMF, at 50 °C<sup>200</sup>. The

overall yield was 79% for the 1-(benzyloxy)pyrazole derivative (**207**) and 70% for the 1-benzyloxy-1,2,3-triazole derivative (**208**). Scheme 25A illustrates the reactions used to obtain this kind of azaxanthenes.

A one-pot approach was also achieved, which represented a faster way of obtaining a pyrazol-containing azaxanthone, albeit with a more modest yield of 25% (Scheme 25B). In this reaction, 1-hydroxypyrazol (**209**) was used as starting material, and was dissolved in THF, at 0 °C, under nitrogen atmosphere, and sodium hydride was added. Afterwards, 2-chlorobenzoyl chloride was added and, after the reaction vessel was cooled to -78 °C, lithium *t*-butyltritylamide was added. Then, the mixture was allowed to heat to room temperature and potassium carbonate was added, followed by heating to 100 °C for 10 hours, yielding the desired azaxanthone (**207**) (Scheme 25B)<sup>200</sup>. The second route aimed to construct the *O*-aryl bond before closing the ring. As such, efforts were placed towards the synthesis of a cyano diaryl-ether (**208**), since the cyano group constitutes an activating group in nucleophilic aromatic substitution (Scheme 25C). 1-Hydroxypyrazol (**209**) reacted with 2-fluorobenzonitrile, in the presence of sodium hydride, in DMF, under nitrogen atmosphere, for three hours, at 70 °C. *n*-Buthyllithium was used for the lithiation, in THF, at -78 °C. Addition of hydrochloric acid led to the desired azaxanthone **207**, with a low yield of approximately 20%. Studies demonstrated that the 4-bromo derivative of the cyano precursor (**209**), which could be obtained with a 99% yield, would give the desired azaxanthone **210** in a more satisfactory 89% yield, only changing the lithiation reagent to LDA, in order to avoid halogen-metal exchange (Scheme 25C)<sup>200</sup>.


Another method towards obtaining five membered ring azaxanthenes was proposed by Singh *et al.*<sup>201</sup>. Starting from a nitro compound **211**, 2-anilino-6-chloro-3-formylchromone (**212**) was formed in dry benzene, with a few drops of acetic acid (Scheme 25D). Reaction with methyl-iodide in the presence of potassium carbonate in acetone yielded 6-chloro-2-(*N*-methylanilino)-3-formylchromone (**213**), which upon reaction with hydrazine

hydrate in aqueous acetonitrile overnight afforded the azaxanthone **214** (Scheme 25D) in 43% yield.


Besides the synthesis of bioactive azaxanthones, the synthesis of thioazaxanthones and their uses as photosensitizers<sup>15, 202</sup> has

been reported as well as the synthesis of more complex pyranoazaxanthones, fused ring azaxanthones<sup>89, 206-218</sup>, and others<sup>203-205</sup>.


#### A. Synthesis of five-membered ring azaxanthone derivatives




#### B. One-pot synthesis of 1-(benzyloxy)pyrazole



#### C. Synthesis of five-membered ring azaxanthone and a halogenated derivative



#### D. Synthesis of a chlorinated five-membered ring azaxanthone derivative



Scheme 25. Other methodologies for the synthesis of xanthones

## REVIEW

## Organic Chemistry Frontiers

## 7. DIVERSIFICATIONS ON THE XANTHONE CORE

Hydroxyl groups are undoubtedly one of the most versatile functional groups when it comes to structural modifications (Scheme 26). A myriad of examples have been described in the literature, including the reaction with alkyl halides for the transformation into alkoxy<sup>43, 59, 69, 219-223</sup>, haloalkoxy<sup>18, 51, 69, 99, 224</sup>, methyl alkyl- or aryl-piperazine moieties<sup>225</sup>, epoxypyranoxy<sup>65, 103, 220-222</sup> oxirane<sup>221</sup>, oxypropanolamines<sup>103</sup>, benzenesulfonamides<sup>226</sup>, propargyloxy<sup>45, 52, 54, 63, 106, 118, 173, 227, 228</sup>. When reacted with other compounds that are not alkyl halides, several other functionalities can be achieved like alkoxy<sup>19, 23, 24, 33, 97</sup> which in certain situations, can be converted into chlorosulfonyl<sup>24</sup> or amine<sup>33</sup>, ethylenedioxy<sup>100</sup>, and allyl<sup>39, 115, 138</sup>. Other transformations include the introduction of glycosyl<sup>55, 110, 132, 135, 229, 230</sup>, ester<sup>17, 62, 227</sup>, amino<sup>151</sup>, and even prenyl<sup>23, 39, 40, 45, 46, 49, 50, 52, 53, 97, 98, 111-114, 117, 130, 133, 221, 231-233</sup> moieties, being the latter further used for the synthesis of caged xanthenes<sup>39, 49, 105, 138, 234</sup> or pyranoxanthenes<sup>40, 45, 46</sup>. Ultimately, hydroxyl groups can be converted into triflates which provide easy access towards amines<sup>227, 235</sup>, imines<sup>116, 140, 235</sup>, aryl<sup>119, 236</sup>, and alkyl xanthenes<sup>155</sup>, or even converted into tosyl groups for further transformations<sup>110</sup> for the introduction of amines. Aminoxanthenes can be obtained through reduction of the nitro group<sup>54, 60, 108, 227, 237, 238</sup>. Further transformation of the amino moiety can be used for the synthesis of oxazol<sup>60</sup>, amide<sup>108, 237-239</sup>, and propargyl<sup>54</sup> xanthone derivatives or even transformed into hydroxyl, chloro, iodo or arylxanthenes via diazonium salts<sup>219</sup>. Carboxyxanthone derivatives can be coupled with amines to obtain the corresponding amides<sup>47, 151</sup> or ester<sup>240</sup>. Albeit methyl group is usually not considered a highly versatile group, after a simple bromination<sup>66, 71, 219, 241</sup> it is possible to obtain several other functionalities. For example, imidazolymethyl<sup>219</sup>, methylhydroxy<sup>66</sup>, methylamine<sup>241</sup> or formyl<sup>66, 71</sup>, being the latter further transformed into methylamine<sup>33, 71</sup>, acid<sup>240</sup>, ester<sup>240</sup> and aldoxime<sup>240</sup> moieties. The presence of a halogen substituent at the xanthone core

also provides an opportunity for further transformations. Recent examples include iodo<sup>137</sup> or bromo<sup>197</sup> to aryl via arylboronic acids, or bromo to alkynyl xanthenes via Sonogashira coupling<sup>197</sup>. On the other hand fluoro can undergo  $S_NAr$  reactions with pyrrolidine and sodium ethoxide<sup>197</sup> or installation of isoprenyl moiety via isoprenyl Grignard reagent<sup>231</sup>.


## Conclusions

The importance of xanthenes and azaxanthenes in medicinal chemistry world led to the employment of advanced synthetic methodologies in these heterocyclic scaffolds. For obtaining xanthenes and azaxanthenes three starting points are generally used: the benzophenone, the diaryl ether or the chromen-4-one.

The benzophenone route consists in a multi-step approach using a benzoic or nicotinic acid derivative, followed by a cyclization, finally leading to the xanthone scaffold. The diaryl ether route focuses on a classic Ullmann condensation between an aryl halide and a phenol derivative. The cyclisation can be achieved by a wide range of methodologies, being also a multi-step approach. On the other hand, the chromen-4-one route is usually a one-step strategy. These reactions are not exclusive to the (aza)xanthone scaffold, being transversal to the synthesis of many heterocycles, such as flavones, coumarins, acridines, among others which expands the scope of the reaction conditions presented herein.

It can also be observed that many approaches are being revisited and recent advances made with the aim of becoming greener methods, such as one-pot reactions, MAOS, solvent- and metal-free strategies. Nanocatalysts represent a significant advancement in catalytic activity over conventional catalytic methods in several synthetic approaches with a diversity of catalysts being employed, such as palladium and magnetically separable nano-CuFe<sub>2</sub>O<sub>4</sub> allowing also high throughputs in purification procedures.

New reagents for transition-metal systems are emerging and include pyridylmagnesium reagents and *N*-heterocyclic carbenes with unique reactivities that have been used in a broad range of organometallic transformations and are expected to increase in the following years. Also, metal-free conditions are being applied such as metal-free variants of the intramolecular Friedel-Crafts acylation, with quaternary ammonium salts like tetrabutylammonium bromide (TBAB)-promoted intramolecular annulation in aqueous medium standing out. Concerning atom economy, one-pot approaches rule with multicomponent and domino/tandem reactions being often applied in cyclization reactions of (aza)xanthenes. Solvent-free conditions and more efficient catalysts are being disclosed such as indium triflate, ytterbium(III) trifluoromethanesulfonate, Cu(0)/Selectfluor system, dichloro(*p*-cymene)ruthenium(II), and also in palladium catalysed carbonylations/C–H activation reactions. Recent trends are the radical cyclization via photoredox catalysis using photosensitizers and the cross dehydrogenative reaction that



Scheme 26. Most common molecular diversifications on substituted xanthenes

involves few steps and can be extended to a wide scope of derivatives.

This review also shows that the xanthone scaffold still gathers a lot of attention not only from a synthetic point of view but due to the diversity of bioactive derivatives being disclosed, the case-studies of this study. New biological applications are being proposed for xanthenes beyond their pharmacological properties with relevance for chemical biology as sensitizing chromophores. Therefore, it is useful to know the vast array of approaches being used, as they can become a valuable toolbox in the search of novel bioactive heterocycles, emphasizing the importance of this privileged scaffold. Moreover, the site-selective late-stage functionalization approaches highlighted herein can guide medicinal chemists in the lead optimization not only of (aza)xanthenes but also of isosters like thioxanthenes, acridones, cromones, flavones, among others. This review can help guide medicinal chemists working with heterocycles by providing chemical tools and reaction conditions to both target and diversity-oriented synthesis approaches.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was supported by national funds through FCT - Foundation for Science and Technology within the scope of UIDB/04423/2020, UIDP/04423/2020, and under the project PTDC/SAU-PUB/28736/2017 (reference POCI-01-0145-FEDER-028736), co-financed by COMPETE 2020, Portugal 2020 and the European Union through the ERDF and by FCT through national funds, as well as CHIRALBIOACTIVE-PI-3RL-IINFACETS-2019. F.D. and M.M. also acknowledge FCT for their PhD grants (SFRH/BD/114681/2019 and SFRH/BD/146211/2019, respectively).

## References

1. G. Wu, T. Zhao, D. Kang, J. Zhang, Y. Song, V. Namasivayam, J. Kongsted, C. Pannecouque, E. De Clercq, V. Poongavanam, X. Liu and P. Zhan, Overview of Recent Strategic Advances in Medicinal Chemistry, *Journal of Medicinal Chemistry*, 2019, **62**, 9375-9414.
2. J. J. Vanden Eynde, A. A. Mangoni, J. Rautio, J. Leprince, Y.-T. Azuma, A. T. García-Sosa, C. Hulme, J. Jampilek, R. Karaman, W. Li, P. A. C. Gomes, D. Hadjipavlou-Litina, R. Capasso, A. Geronikaki, L. Cerchia, J.-M. Sabatier, R. Ragno, T. Tuccinardi, A. Trabocchi, J.-Y. Winum, F. J. Luque, K. Prokai-Tatrai, M. Spetea, M. Gütschow, I. Kosalec, C. Guillou, M. H. Vasconcelos, G. Kokotos, G. Rastelli, M. E. de Sousa, C. Manera, S. Gemma, S. Mangani, C. Siciliano, S. Galdiero, H. Liu, P. J. H. Scott, C. de los Ríos, L. A. Agrofoglio, S. Collina, R. C. Guedes and D. Muñoz-Torrero, Breakthroughs in Medicinal Chemistry: New Targets and Mechanisms, New Drugs, New Hopes-6, *Molecules*, 2019, **25**, 119.
3. A. P. Taylor, R. P. Robinson, Y. M. Fobian, D. C. Blakemore, L. H. Jones and O. Fadeyi, Modern advances in heterocyclic chemistry in drug discovery, *Organic & Biomolecular Chemistry*, 2016, **14**, 6611-6637.
4. J. Jampilek, Heterocycles in Medicinal Chemistry, *Molecules*, 2019, **24**, 3839.
5. Z. Feng, X. Lu, L. Gan, Q. Zhang and L. Lin, Xanthenes, A Promising Anti-Inflammatory Scaffold: Structure, Activity, and Drug Likeness Analysis, *Molecules*, 2020, **25**, 598.
6. Z. Salman, J. Yu-Qing, L. Bin, P. Cai-Yun, C. M. Iqbal, R. Attaur and W. Wei, Antioxidant Nature Adds Further Therapeutic Value: An Updated Review on Natural Xanthenes and Their Glycosides, *Digital Chinese Medicine*, 2019, **2**, 166-192.
7. R. E.-S. Hesham, M. H. E.-G. Dina, A. E.-B. Mai, F. Z. Mervat, G. Ulf, L. Sonny and V. Rob, Naturally Occurring Xanthenes; Latest Investigations: Isolation, Structure Elucidation and Chemosystematic Significance, *Current Medicinal Chemistry*, 2009, **16**, 2581-2626.
8. D. R. P. Loureiro, J. X. Soares, J. C. Costa, Á. F. Magalhães, C. M. G. Azevedo, M. M. M. Pinto and C. M. M. Afonso, Structures, Activities and Drug-Likeness of Anti-Infective Xanthone Derivatives Isolated from the Marine Environment: A Review, *Molecules*, 2019, **24**, 243.
9. M. E. Sousa and M. M. M. Pinto, Synthesis of Xanthenes: An Overview, *Current Medicinal Chemistry*, 2005, **12**, 2447-2479.
10. A. Carlos Miguel Goncalves, A. Carlos Manuel Magalhaes and P. Madalena Maria Magalhaes, Routes to Xanthenes: An Update on the Synthetic Approaches, *Current Organic Chemistry*, 2012, **16**, 2818-2867.
11. A. S. Gomes, P. Brandão, C. S. G. Fernandes, M. R. P. C. d. Silva, M. E. d. S. P. d. Sousa and M. M. d. M. Pinto, Drug-like Properties and ADME of Xanthone Derivatives: The Antechamber of Clinical Trials, *Current Medicinal Chemistry*, 2016, **23**, 3654-3686.
12. Á. Santos, J. X. Soares, S. Cravo, M. E. Tiritan, S. Reis, C. Afonso, C. Fernandes and M. M. M. Pinto, Lipophilicity assessment in drug discovery: Experimental and theoretical methods applied to xanthone derivatives, *Journal of Chromatography B*, 2018, **1072**, 182-192.
13. E. Vitaku, D. T. Smith and J. T. Njardarson, Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals, *Journal of Medicinal Chemistry*, 2014, **57**, 10257-10274.
14. J. Yan, M. Cheng, F. Hu and Y. Hu, Direct synthesis of functional azaxanthenes by using a domino three-component reaction, *Organic Letters*, 2012, **14**, 3206-3209.
15. P. Atkinson, K. S. Findlay, F. Kielar, R. Pal, D. Parker, R. A. Poole, H. Puschmann, S. L. Richardson, P. A. Stenson, A. L. Thompson and J. Yu, Azaxanthenes and azathioxanthenes are effective sensitizers for europium and terbium luminescence, *Organic & Biomolecular Chemistry*, 2006, **4**, 1707-1722.
16. M. M. M. Pinto, M. E. Sousa and M. S. J. Nascimento, Xanthone Derivatives: New Insights in Biological Activities, *Current Medicinal Chemistry*, 2005, **12**, 2517-2538.
17. X. Chen, J. Leng, K. P. Rakesh, N. Darshini, T. Shubhavathi, H. K. Vivek, N. Mallesha and H. L. Qin, Synthesis and molecular docking studies of xanthone attached amino



REVIEW

Organic Chemistry Frontiers

- acids as potential antimicrobial and anti-inflammatory agents, *Medchemcomm*, 2017, **8**, 1706-1719.
18. A. Das, M. M. Shaikh and S. Jana, Design, synthesis, and in vitro antibacterial screening of some novel 3-pentyloxy-1-hydroxyxanthone derivatives, *Medicinal Chemistry Research*, 2014, **23**, 436-444.
19. T. Kodama, T. Ito, D. F. Dibwe, S. Y. Woo and H. Morita, Syntheses of benzophenone-xanthone hybrid polyketides and their antibacterial activities, *Bioorganic & Medicinal Chemistry Letters*, 2017, **27**, 2397-2400.
20. J. Araujo, C. Fernandes, M. Pinto and M. E. Tiritan, Chiral derivatives of Xanthenes with antimicrobial activity, *Molecules*, 2019, **24**, 314/311-314/329.
21. A. K. Kumar, V. Sunitha, B. Shankar, T. M. Krishna, C. A. Lincoln and P. Jalapathi, Synthesis and antimicrobial screening of novel 9-(2- (1H-1,2,3-triazol-4-yl)methoxy phenyl)-3,3,6,6-tetramethyl-3,4,5,6 ,7,9-hexahydro-1H-xanthene-1,8(2H)-dione derivatives, *Russian Journal of General Chemistry*, 2017, **87**, 2011-2020.
22. G. Liao, J. Fan, L. Ludwig-Radtke, K. Backhaus and S. M. Li, Increasing Structural Diversity of Natural Products by Michael Addition with ortho-Quinone Methide as the Acceptor, *Journal of Organic Chemistry*, 2020, **85**, 1298-1307.
23. S. M. Lin, J. J. Koh, T. T. Aung, F. H. Lim, J. G. Li, H. X. Zou, L. Wang, R. Lakshminarayanan, C. Verma, Y. J. Wang, D. T. H. Tan, D. R. Cao, R. W. Beuerman, L. Ren and S. P. Liu, Symmetrically Substituted Xanthone Amphiphiles Combat Gram-Positive Bacterial Resistance with Enhanced Membrane Selectivity, *Journal of Medicinal Chemistry*, 2017, **60**, 1362-1378.
24. M. Malekpoor, S. Gharaghani, A. Sharifzadeh, S. N. Mirsattari and A. R. Massah, Synthesis and antibacterial evaluation of novel xanthone sulfonamides, *Journal of Chemical Research*, 2015, DOI: 10.3184/174751915x14373971129805, 433-437.
25. A. C. Micheletti, N. K. Honda, N. C. P. Carvalho, D. P. de Lima and A. Beatriz, Design, Synthesis and in vitro Antimicrobial Activity Evaluation of Novel Hybrids of Lichexantone-THC Derivatives, *Orbital-the Electronic Journal of Chemistry*, 2015, **7**, 301-307.
26. H. X. Zou, J. J. Koh, J. G. Li, S. X. Qiu, T. T. Aung, H. F. Lin, R. Lakshminarayanan, X. P. Dai, C. Tang, F. H. Lim, L. Zhou, A. L. Tan, C. Verma, D. T. H. Tan, H. S. O. Chan, P. Saraswathi, D. R. Cao, S. P. Liu and R. W. Beuerman, Design and Synthesis of Amphiphilic Xanthone-Based, Membrane-Targeting Antimicrobials with Improved Membrane Selectivity (vol 56, pg 2359, 2013), *Journal of Medicinal Chemistry*, 2013, **56**, 3747-3747.
27. L. J. Bessa, A. Palmeira, A. S. Gomes, V. Vasconcelos, E. Sousa, M. Pinto and P. M. d. Costa, Synergistic Effects Between Thioxanthenes and Oxacillin Against Methicillin-Resistant *Staphylococcus aureus*, *Microbial Drug Resistance*, 2015, **21**, 404-415.
28. D. Resende, P. Pereira-Terra, A. S. Inacio, P. M. da Costa, E. Pinto, E. Sousa and M. M. M. Pinto, Lichen Xanthenes as Models for New Antifungal Agents, *Molecules*, 2018, **23**.
29. E. Pinto, C. Afonso, S. Duarte, L. Vale-Silva, E. Costa, E. Sousa and M. Pinto, Antifungal Activity of Xanthenes: Evaluation of their Effect on Ergosterol Biosynthesis by High-performance Liquid Chromatography, *Chemical Biology & Drug Design*, 2011, **77**, 212-222.
30. A. Pedro Gonçalves, N. Silva, C. Oliveira, D. J. Kowbel, N. L. Glass, A. Kijjoo, A. Palmeira, E. Sousa, M. Pinto and A. Videira, Transcription profiling of the *Neurospora crassa* response to a group of synthetic (thio)xanthenes and a natural acetophenone, *Genom Data*, 2015, **4**, 26-32.
31. R. T. Lima, H. Seca, A. Palmeira, M. X. Fernandes, F. Castro, M. Correia-da-Silva, M. S. J. Nascimento, E. Sousa, M. Pinto and M. H. Vasconcelos, Sulfated Small Molecules Targeting EBV in Burkitt Lymphoma: From In Silico Screening to the Evidence of In Vitro Effect on Viral Episomal DNA, *Chemical Biology & Drug Design*, 2013, **81**, 631-644.
32. E. V. Buravlev, O. G. Shevchenko, A. A. Anisimov and K. Y. Suponitsky, Novel Mannich bases of alpha- and gamma-mangostins: Synthesis and evaluation of antioxidant and membrane-protective activity, *European Journal of Medicinal Chemistry*, 2018, **152**, 10-20.
33. I. Cruz, P. Puthongking, S. Cravo, A. Palmeira, H. Cidade, M. Pinto and E. Sousa, Xanthone and Flavone Derivatives as Dual Agents with Acetylcholinesterase Inhibition and Antioxidant Activity as Potential Anti-Alzheimer Agents, *Journal of Chemistry*, 2017, DOI: 10.1155/2017/8587260.
34. J. M. Khurana, A. Chaudhary, A. Lumb and B. Nand, Efficient one-pot syntheses of dibenzo a,i xanthene-diones and evaluation of their antioxidant activity, *Canadian Journal of Chemistry-Revue Canadienne De Chimie*, 2012, **90**, 739-746.
35. E. Yuanita, H. D. Pranowo, D. Siswanta, R. T. Swasono, Mustofa, A. K. Zulkarnain, J. Syahri and Jumina, ONE-POT SYNTHESIS, ANTIOXIDANT ACTIVITY AND TOXICITY EVALUATION OF SOME HYDROXYXANTHONES, *Chemistry & Chemical Technology*, 2018, **12**, 290-295.
36. B. D. Zhou, L. L. Zeng, Y. G. Tong, J. Y. Fang, Z. P. Ruan, X. Y. Zeng, Y. Y. Fang, G. F. Xu and D. B. Hu, Synthesis and antitumor, antityrosinase, and antioxidant activities of xanthone (vol 20, pg 467, 2018), *Journal of Asian Natural Products Research*, 2019, **21**, 688-688.
37. H. Cidade, V. Rocha, A. Palmeira, C. Marques, M. E. Tiritan, H. Ferreira, J. S. Lobo, I. F. Almeida, M. E. Sousa and M. Pinto, In silico and in vitro antioxidant and cytotoxicity evaluation of oxygenated xanthone derivatives, *Arabian Journal of Chemistry*, 2020, **13**, 17-26.
38. R. Urbatzka, S. Freitas, A. Palmeira, T. Almeida, J. Moreira, C. Azevedo, C. Afonso, M. Correia-da-Silva, E. Sousa, M. Pinto and V. Vasconcelos, Lipid reducing activity and toxicity profiles of a library of polyphenol derivatives, *European Journal of Medicinal Chemistry*, 2018, **151**, 272-284.
39. O. Chantarasriwong, A. T. Milcarek, T. H. Morales, A. L. Settle, C. O. Rezende, Jr., B. D. Althufairi, M. A. Theodoraki, M. L. Alpaugh and E. A. Theodorakis, Synthesis, structure-activity relationship and in vitro pharmacodynamics of A-ring modified caged xanthenes in a preclinical model of inflammatory breast cancer, *European Journal of Medicinal Chemistry*, 2019, **168**, 405-413.
40. C. T. Yen, K. Nakagawa-Goto, T. L. Hwang, S. L. Morris-Natschke, K. F. Bastow, Y. C. Wu and K. H. Lee, Design and synthesis of gambogic acid analogs as potent cytotoxic and anti-inflammatory agents, *Bioorganic & Medicinal Chemistry Letters*, 2012, **22**, 4018-4022.
41. D. Zelaszczyk, A. Lipkowska, N. Szkaradek, K. Sloczynska, A. Gunia-Krzyzak, T. Librowski and H. Marona, Synthesis and preliminary anti-inflammatory evaluation of xanthone

- derivatives, *Heterocyclic Communications*, 2018, **24**, 231-236.
42. M. Correia-da-Silva, E. Sousa, B. Duarte, F. Marques, F. Carvalho, L. M. Cunha-Ribeiro and M. M. M. Pinto, Polysulfated Xanthenes: Multipathway Development of a New Generation of Dual Anticoagulant/Antiplatelet Agents, *Journal of Medicinal Chemistry*, 2011, **54**, 5373-5384.
  43. R. Shen, W. H. Wang and G. L. Yang, DNA Binding Property and Antitumor Evaluation of Xanthone with Dimethylamine Side Chain, *Journal of Fluorescence*, 2014, **24**, 959-966.
  44. F. M. Soliman, M. M. Said, M. Youns and S. A. Darwish, Chemistry of phosphorus ylides 32: synthesis of phosphoranylidene-pyrano- and -cyclobutyl-xanthenones with potential antitumor activity, *Monatshefte Fur Chemie*, 2012, **143**, 965-973.
  45. C. M. G. Azevedo, C. M. M. Afonso, J. X. Soares, S. Reis, D. Sousa, R. T. Lima, M. H. Vasconcelos, M. Pedro, J. Barbosa, L. Gales and M. M. M. Pinto, Pyranoxanthenes: Synthesis, growth inhibitory activity on human tumor cell lines and determination of their lipophilicity in two membrane models, *European Journal of Medicinal Chemistry*, 2013, **69**, 798-816.
  46. C. M. G. Azevedo, C. M. M. Afonso, D. Sousa, R. T. Lima, M. H. Vasconcelos, M. Pedro, J. Barbosa, A. G. Correa, S. Reis and M. M. M. Pinto, Multidimensional optimization of promising antitumor xanthone derivatives, *Bioorganic & Medicinal Chemistry*, 2013, **21**, 2941-2959.
  47. C. Fernandes, K. Masawang, M. E. Tiritan, E. Sousa, V. de Lima, C. Afonso, H. Bousbaa, W. Sudprasert, M. Pedro and M. M. Pinto, New chiral derivatives of xanthenes: Synthesis and investigation of enantioselectivity as inhibitors of growth of human tumor cell lines, *Bioorganic & Medicinal Chemistry*, 2014, **22**, 1049-1062.
  48. W. L. Lan, J. C. Wei, J. K. Qin, Z. M. Yang, G. F. Su and Z. K. Dai, Synthesis and Biological Evaluation of 1, 3-Dihydroxyxanthone Mannich Base Derivatives as Potential Antitumor Agents, *Letters in Drug Design & Discovery*, 2013, **10**, 689-698.
  49. X. Li, X. J. Zhang, X. J. Wang, N. G. Li, C. J. Lin, Y. Gao, Z. Q. Yu, Q. L. Guo and Q. D. You, Synthesis and Anti-tumor Evaluation of B-ring Modified Caged Xanthone Analogues of Gambogic Acid, *Chinese Journal of Chemistry*, 2012, **30**, 35-42.
  50. C. K. Lim, L. Y. Tho, Y. M. Lim, S. A. A. Shah and J. F. F. Weber, Synthesis of 1,3,6-Trioxxygenated Prenylated Xanthone Derivatives as Potential Antitumor Agents, *Letters in Organic Chemistry*, 2012, **9**, 549-555.
  51. L. Luo, J. K. Qin, Z. K. Dai and S. H. Gao, Synthesis and biological evaluation of novel benzo b xanthone derivatives as potential antitumor agents, *Journal of the Serbian Chemical Society*, 2013, **78**, 1301-1308.
  52. X. J. Zhan, X. Li, H. P. Sun, X. J. Wang, L. Zhao, Y. Gao, X. R. Liu, S. L. Zhang, Y. Y. Wang, Y. R. Yang, S. Zeng, Q. L. Guo and Q. D. You, Garcinia Xanthenes as Orally Active Antitumor Agents, *Journal of Medicinal Chemistry*, 2013, **56**, 276-292.
  53. X. J. Zhang, X. Li, S. F. Ye, Y. Zhang, L. Tao, Y. Gao, D. D. Gong, M. Y. Xi, H. Y. Meng, M. Q. Zhang, W. L. Gao, X. L. Xu, Q. L. Guo and Q. D. You, Synthesis, SAR and Biological Evaluation of Natural and Non-natural Hydroxylated and Prenylated Xanthenes as Antitumor Agents, *Medicinal Chemistry*, 2012, **8**, 1012-1025.
  54. Y. Zou, Q. J. Zhao, H. G. Hu, L. N. Hu, S. C. Yu, M. J. Xu and Q. Y. Wu, Synthesis and in vitro antitumor activities of xanthone derivatives containing 1,4-disubstituted-1,2,3-triazole moiety, *Archives of Pharmacol Research*, 2012, **35**, 2093-2104.
  55. A. Alves, J. Campos, D. Ferreira, P. C. Costa, A. Alves, M. Correia-da-Silva, E. Sousa, M. Pinto, M. Correia-da-Silva, E. Sousa, H. Bousbaa, M. Pinto, C. Nunes, P. M. A. Silva, H. Bousbaa and F. Rodrigues, Discovery of a New Xanthone against Glioma: Synthesis and Development of (Pro)liposome Formulations, *Molecules*, 2019, **24**.
  56. J. Barbosa, R. T. Lima, D. Sousa, A. S. Gomes, A. Palmeira, H. Seca, K. Choosang, P. Pakkong, H. Bousbaa, M. M. Pinto, E. Sousa, M. H. Vasconcelos and M. Pedro, Screening a Small Library of Xanthenes for Antitumor Activity and Identification of a Hit Compound which Induces Apoptosis, *Molecules*, 2016, **21**, 81.
  57. C. Elisangela, S. Emilia, N. Nair, S. J. N. Maria and M. M. P. Madalena, Synthesis of Xanthenes and Benzophenones as Inhibitors of Tumor Cell Growth, *Letters in Drug Design & Discovery*, 2010, **7**, 487-493.
  58. A. Palmeira, A. Paiva, E. Sousa, H. Seca, G. M. Almeida, R. T. Lima, M. X. Fernandes, M. Pinto and M. H. Vasconcelos, Insights into the In Vitro Antitumor Mechanism of Action of a New Pyranoxanthone, *Chemical Biology & Drug Design*, 2010, **76**, 43-58.
  59. P. S. Bairy, A. Das, L. M. Nainwal, T. K. Mohanta, M. K. Kumawat, P. K. Mohapatra and P. Parida, Design, synthesis and anti-diabetic activity of some novel xanthone derivatives targeting alpha-glucosidase, *Bangladesh Journal of Pharmacology*, 2016, **11**, 308-318.
  60. S. M. Ding, T. Lan, G. J. Ye, J. J. Huang, Y. Hu, Y. R. Zhu and B. Wang, Novel oxazolxanthone derivatives as a new type of alpha-glucosidase inhibitor: synthesis, activities, inhibitory modes and synergetic effect, *Bioorganic & Medicinal Chemistry*, 2018, **26**, 3370-3378.
  61. Z. Heydari, M. Mohammadi-Khanaposhtani, S. Imanparast, M. A. Faramarzi, M. Mandavi, P. R. Ranjbar and B. Larijani, Pyrano 3,2-c quinoline Derivatives as New Class of alpha-glucosidase Inhibitors to Treat Type 2 Diabetes: Synthesis, in vitro Biological Evaluation and Kinetic Study, *Medicinal Chemistry*, 2019, **15**, 8-16.
  62. G. L. Li, C. Y. Cai, J. Y. He, L. Rao, L. Ma, Y. Liu and B. Wang, Synthesis of 3-acyloxyxanthone derivatives as alpha-glucosidase inhibitors: A further insight into the 3-substituents' effect, *Bioorganic & Medicinal Chemistry*, 2016, **24**, 1431-1438.
  63. G.-J. Ye, T. Lan, Z.-X. Huang, X.-N. Cheng, C.-Y. Cai, S.-M. Ding, M.-L. Xie and B. Wang, Design and synthesis of novel xanthone-triazole derivatives as potential antidiabetic agents:  $\alpha$ -Glucosidase inhibition and glucose uptake promotion, *European Journal of Medicinal Chemistry*, 2019, **177**, 362-373.
  64. P. F. Cheng, L. J. Zhu, W. Guo, W. F. Liu, J. Z. Yao, G. Q. Dong, Y. Q. Zhang, C. L. Zhuang, C. Q. Sheng, Z. Y. Miao and W. N. Zhang, Synthesis of novel benzoxanthone analogues as non-Camptothecin topoisomerase I inhibitors, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2012, **27**, 437-442.

REVIEW

Organic Chemistry Frontiers

65. S. Park, E. Hong, S. Y. Kwak, K. Y. Jun, E. S. Lee, Y. Kwon and Y. Na, Synthesis and biological evaluation of C1-O-substituted-3-(3-butylamino-2-hydroxy-propoxy)-xanthen-9-one as topoisomerase II  $\alpha$  catalytic inhibitors, *European Journal of Medicinal Chemistry*, 2016, **123**, 211-225.
66. E. Martins, V. Silva, A. Lemos, A. Palmeira, P. Puthongking, E. Sousa, C. Rocha-Pereira, C. I. Ghanem, H. Carmo, F. Remiao and R. Silva, Newly synthesized oxygenated xanthenes as potential P-glycoprotein activators: in vitro, ex vivo, and in silico studies, *Molecules*, 2019, **24**, 707.
67. R. Silva, E. Sousa, H. Carmo, A. Palmeira, D. J. Barbosa, M. Gameiro, M. Pinto, M. de Lourdes Bastos and F. Remião, Induction and activation of P-glycoprotein by dihydroxylated xanthenes protect against the cytotoxicity of the P-glycoprotein substrate paraquat, *Archives of Toxicology*, 2014, **88**, 937-951.
68. E. Sousa, A. Palmeira, A. S. Cordeiro, B. Sarmiento, D. Ferreira, R. T. Lima, M. Helena Vasconcelos and M. Pinto, Bioactive xanthenes with effect on P-glycoprotein and prediction of intestinal absorption, *Medicinal Chemistry Research*, 2013, **22**, 2115-2123.
69. M. S. Alawi, T. A. Awad, M. A. Mohamed, A. Khalid, E. M. O. Ismail, F. Alfatih, S. Naz and Z. Ul-Haq, Insights into the molecular basis of acetylcholinesterase inhibition by xanthenes: an integrative in silico and in vitro approach, *Molecular Simulation*, 2020, **46**, 253-261.
70. C. A. Menendez, B. Biscussi, S. Accordino, A. P. Murray, D. C. Gerbino and G. A. Appignanesi, Design, synthesis and biological evaluation of 1,3-dihydroxyxanthone derivatives: Effective agents against acetylcholinesterase, *Bioorganic Chemistry*, 2017, **75**, 201-209.
71. A. Lemos, A. S. Gomes, J. B. Loureiro, P. Brandao, A. Palmeira, M. M. M. Pinto, L. Saraiva and M. E. Sousa, Synthesis, biological evaluation, and in silico studies of novel aminated xanthenes as potential p53-activating agents, *Molecules*, 2019, **24**, 1975pp.
72. S. Gomes, L. Raimundo, J. Soares, J. B. Loureiro, M. Leão, H. Ramos, M. N. Monteiro, A. Lemos, J. Moreira, M. Pinto, P. Chlapek, R. Veselska, E. Sousa and L. Saraiva, New inhibitor of the TAp73 interaction with MDM2 and mutant p53 with promising antitumor activity against neuroblastoma, *Cancer Letters*, 2019, **446**, 90-102.
73. M. Leão, S. Gomes, J. Pedraza-Chaverri, N. Machado, E. Sousa, M. Pinto, A. Inga, C. Pereira and L. Saraiva,  $\alpha$ -Mangostin and Gambogic Acid as Potential Inhibitors of the p53-MDM2 Interaction Revealed by a Yeast Approach, *Journal of Natural Products*, 2013, **76**, 774-778.
74. M. Leão, C. Pereira, A. Bisio, Y. Ciribilli, A. M. Paiva, N. Machado, A. Palmeira, M. X. Fernandes, E. Sousa, M. Pinto, A. Inga and L. Saraiva, Discovery of a new small-molecule inhibitor of p53-MDM2 interaction using a yeast-based approach, *Biochemical Pharmacology*, 2013, **85**, 1234-1245.
75. US 7,772,255 B2, 2010.
76. G. Kolokythas, I. K. Kostakis, N. Pouli, P. Marakos, D. Kletsas and H. Pratsinis, Synthesis and cytotoxic activity of some new azapyranoxanthenone aminoderivatives, *Bioorganic & Medicinal Chemistry*, 2003, **11**, 4591-4598.
77. EP 0 075 607 B1, 1985.
78. F. J. Villani, T. A. Mann, E. A. Wefer, J. Hannon, L. L. Larca, M. J. Landon, W. Spivak, D. Vashi and S. Tozzi, Benzopyranopyridine derivatives. 1. Aminoalkyl derivatives of the azaxanthenes as bronchodilating agents, *Journal of Medicinal Chemistry*, 1975, **18**, 1-8.
79. W. Meng, Y. Dong, J. Liu, Z. Wang, X. Zhong, R. Chen, H. Zhou, M. Lin, L. Jiang, F. Gao, T. Xu, Q. Chen and X. Zeng, A clinical evaluation of amlexanox oral adhesive pellicles in the treatment of recurrent aphthous stomatitis and comparison with amlexanox oral tablets: a randomized, placebo controlled, blinded, multicenter clinical trial, *Trials*, 2009, **10**, 30-30.
80. D. Rakel, in *Integrative Medicine (Fourth Edition)*, ed. D. Rakel, Elsevier, 2018, DOI: <https://doi.org/10.1016/B978-0-323-35868-2.00075-X>, pp. 747-752.e741.
81. A. Michael, On the action of aromatic oxy-acids on phenols, *Berichte der Deutschen Chemischen Gesellschaft* 1883, **5**, 81.
82. S. v. Kostanecki and B. Nessler, Synthesen von Oxyxanthonen, *Berichte der deutschen chemischen Gesellschaft*, 1891, **24**, 1894-1897.
83. F. M. Dean, in *The Total Synthesis of Natural Products*, ed. J. Apsimon, 1973, pp. 534-562.
84. M. Afzal and J. M. Al-Hassan, Synthesis and Biosynthesis of Phyttoxanthenes, *Heterocycles*, 1980, **14**, 1173-1205.
85. J. Ribeiro, C. Veloso, C. Fernandes, M. E. Tiritan and M. M. M. Pinto, Carboxyxanthenes: bioactive agents and molecular scaffold for synthesis of analogues and derivatives, *Molecules*, 2019, **24**, 180/181-180/133.
86. R. Khanna, A. Dalal, R. Kumar and R. C. Kamboj, Synthesis of Xanthenones: A Review, *Chemistryselect*, 2016, **1**, 840-851.
87. C. M. M. Santos, D. Pinto, V. L. M. Silva and A. M. S. Silva, Arylxanthenes and arylacridones: a synthetic overview, *Pure and Applied Chemistry*, 2016, **88**, 579-594.
88. A. M. Paiva, M. M. Pinto and E. Sousa, A Century of Thioxanthenes: Through Synthesis and Biological Applications, *Current Medicinal Chemistry*, 2013, **20**, 2438-2457.
89. D. Parker, J. W. Walton, L. Lamarque and J. M. Zwier, Comparative Study of the Luminescence Properties and Relative Stability of a Series of Europium(III) Complexes Bearing One to Four Coordinated Azaxanthone Groups, *European Journal of Inorganic Chemistry*, 2010, **2010**, 3961-3966.
90. C. Coenjarts and J. C. Scaiano, Reaction Pathways Involved in the Quenching of the Photoactivated Aromatic Ketones Xanthone and 1-Azaxanthone by Polyalkylbenzenes, *Journal of the American Chemical Society*, 2000, **122**, 3635-3641.
91. S. Corrent, L. J. Martínez, J. C. Scaiano, H. García and V. Fornés, Intrazeolite Photochemistry. Photochemistry of 1-Azaxanthone in Zeolites in the Presence of Hydrogen Donors, Electron Donors, and Energy Acceptors, *The Journal of Physical Chemistry B*, 1999, **103**, 8097-8103.
92. L. J. Martínez and J. C. Scaiano, The Photochemistry of 1-Azaxanthone in Aqueous Solutions and in Micellar Environments, *The Journal of Physical Chemistry A*, 1999, **103**, 203-208.
93. M. Sakamoto, X. Cai, M. Hara, S. Tojo, M. Fujitsuka and T. Majima, Anomalous Fluorescence from the Azaxanthone Ketyl Radical in the Excited State, *Journal of the American Chemical Society*, 2005, **127**, 3702-3703.

94. J. C. Scaiano, D. Weldon, C. N. Pliva and L. J. Martínez, Photochemistry and Photophysics of 1-Azaxanthone in Organic Solvents, *The Journal of Physical Chemistry A*, 1998, **102**, 6898-6903.
95. P. K. Grover, G. D. Shah and R. C. Shah, Xanthon. Part IV. A new synthesis of hydroxyxanthon. and hydroxybenzophenones, *Journal of the Chemical Society (Resumed)*, 1955, DOI: 10.1039/JR9550003982, 3982-3985.
96. Gover, G. D. Shah and R. C. Shah, Xanthon. Part III- A New Synthesis of Hydroxyxanthon. and Hydroxybenzophenones. Part 111, *Journal of Scientific and Industrial Research, India*, 1954, **13**, 175.
97. J. K. Qin, W. L. Lan, Z. Liu, J. Huang, H. Tang and H. S. Wang, Synthesis and biological evaluation of 1, 3-dihydroxyxanthone mannich base derivatives as anticholinesterase agents, *Chemistry Central Journal*, 2013, **7**.
98. Q. R. Shen, Y. Dai, G. H. Wang, F. Yao, Y. H. Duan, H. F. Chen, W. G. Zhang, X. K. Zhang and X. S. Yao, Total synthesis and RXR  $\alpha$ -mediated transcription studies of neriifolone B and related compounds, *Bioorganic & Medicinal Chemistry*, 2014, **22**, 2671-2677.
99. Z. M. Yang, J. Huang, J. K. Qin, Z. K. Dai, W. L. Lan, G. F. Su, H. Tang and F. Yang, Design, synthesis and biological evaluation of novel 1-hydroxyl-3-aminoalkoxy xanthone derivatives as potent anticancer agents, *European Journal of Medicinal Chemistry*, 2014, **85**, 487-497.
100. N. Guo, J. Liu, L. Qin, D. Jiang, X. You, K. Lu, Y. O. Teng and P. Yu, Synthesis and antitumor activity evaluation of a novel series of xanthone derivatives, *Journal of Asian Natural Products Research*, 2015, **17**, 377-383.
101. J. Liu, J. R. Zhang, H. L. Wang, Z. J. Liu, C. Zhang, Z. L. Jiang and H. R. Chen, Synthesis of xanthone derivatives and studies on the inhibition against cancer cells growth and synergistic combinations of them, *European Journal of Medicinal Chemistry*, 2017, **133**, 50-61.
102. Z. L. Jiang, J. Liu, F. Zhou, J. R. Zhang, Z. J. Liu, C. Zhang and H. R. Chen, One-step Construction of Xanthone Scaffold Assisted by Microwave Irradiation to Optimize the Synthesis of DMXAA, *Chemical Research in Chinese Universities*, 2018, **34**, 918-922.
103. O. Goshain and B. Ahmed, Antihypertensive activity, toxicity and molecular docking study of newly synthesized xanthone derivatives (xanthonypropanolamine), *PLoS One*, 2019, **14**, e0220920.
104. J. Liu, H. Bao, H. Wang, Q. Luo, J. Zuo, Z. Liu, S. Qiu, X. Sun and X. Liu, Synthesis of xanthone derivatives and anti-hepatocellular carcinoma potency evaluation: induced apoptosis, *RSC Adv.*, 2019, **9**, 40781-40791.
105. X. J. Zhang, X. Li, H. P. Sun, Z. Y. Jiang, L. Tao, Y. Gao, Q. L. Guo and Q. D. You, Synthesis and evaluation of novel azacaged Garcinia xanthon. derivatives, *Organic & Biomolecular Chemistry*, 2012, **10**, 3288-3299.
106. M. Dai, X. Yuan, Z. J. Zhu, L. Shan, R. H. Liu, Q. Y. Sun and W. D. Zhang, Efficient Total Synthesis and Biological Activities of 6-Deoxyisocareub. in, *Archiv Der Pharmazie*, 2013, **346**, 314-320.
107. S. S. Mao, H. H. Zheng, C. T. Luo and H. R. Chen, Concise Syntheses of 1,3-Dihydroxyxanthon. derivatives and Their Clearance Effects against Nitrite, *Chinese Journal of Organic Chemistry*, 2014, **34**, 515-521.
108. F. L. Qu, G. J. Bai, W. S. Dong, Y. Yang, Y. S. Jin, Q. G. Meng, Q. Y. Wu, W. Guo and S. C. Yu, Synthesis and Evaluation of Novel Xanthone Derivatives as Potent AChE Inhibitors, *Asian Journal of Chemistry*, 2014, **26**, 3496-3498.
109. X. Li, Y. Zou, Q. J. Zhao, Y. Yang, M. C. Wu, T. Huang, H. G. Hu and Q. Y. Wu, Synthesis, Biological Evaluation, and Molecular Docking Studies of Xanthone Sulfonamides as ACAT Inhibitors, *Chemical Biology & Drug Design*, 2015, **85**, 394-403.
110. G. P. Song, S. M. Li, H. Z. Si, Y. B. Li, Y. S. Li, J. H. Fan, Q. Q. Liang, H. B. He, H. M. Ye and Z. N. Cui, Synthesis and bioactivity of novel xanthone and thioxanthone L-rhamnopyranosides, *Rsc Advances*, 2015, **5**, 36092-36103.
111. J. J. Koh, H. X. Zou, S. M. Lin, H. F. Lin, R. T. Soh, F. H. Lim, W. L. Koh, J. G. Li, R. Lakshminarayanan, C. Verma, D. T. H. Tan, D. R. Cao, R. W. Beuerman and S. P. Liu, Nonpeptidic Amphiphilic Xanthone Derivatives: Structure-Activity Relationship and Membrane-Targeting Properties, *Journal of Medicinal Chemistry*, 2016, **59**, 171-193.
112. K. D. Reichl, M. J. Smith, M. K. Song, R. P. Johnson and J. A. Porco, Biomimetic Total Synthesis of (+/-)-Griffithxanthone via a Cationic Cycloaddition-Cyclization Cascade, *Journal of the American Chemical Society*, 2017, **139**, 14053-14056.
113. J. Syahri, E. Yuanita, B. A. Nurohmah, M. H. Wathon, R. Syafri, R. Armunanto and B. Purwono, Xanthone as Antimalarial: QSAR Analysis, Synthesis, Molecular Docking and In-vitro Antimalarial Evaluation, *Oriental Journal of Chemistry*, 2017, **33**, 29-40.
114. X. Q. Chi, C. T. Zi, H. M. Li, L. Yang, Y. F. Lv, J. Y. Li, B. Hou, F. C. Ren, J. M. Hu and J. Zhou, Design, synthesis and structure-activity relationships of mangostin analogs as cytotoxic agents, *Rsc Advances*, 2018, **8**, 41377-41388.
115. T. Khammee, W. Jongsu, M. Kuno and S. Suksanrarn, Allylxanthone Derivatives as Xanthine Oxidase Inhibitors: Synthesis, Biological Evaluation and Molecular Docking Study, *Oriental Journal of Chemistry*, 2018, **34**, 38-44.
116. P. H. Wang, L. L. Jiang, Y. Cao, D. Y. Ye and L. Zhou, The Design and Synthesis of N-Xanthone Benzenesulfonamides as Novel Phosphoglycerate Mutase 1 (PGAM1) Inhibitors, *Molecules*, 2018, **23**.
117. P. F. Zhou, A. J. Hou and Y. Wang, Total Synthesis of Cudratricusxanthone B, *Chinese Journal of Organic Chemistry*, 2018, **38**, 156-161.
118. J. Wu, J. Dai, Y. Zhang, J. Wang, L. Huang, H. Ding, T. Li, Y. Zhang, S. Yu, J. Wu, J. Dai, Y. Zhang, T. Li, Y. Zhang and J. Mao, Synthesis of Novel Xanthone Analogues and Their Growth Inhibitory Activity Against Human Lung Cancer A549 Cells, *Drug Des Devel Ther*, 2019, **13**, 4239-4246.
119. L. Yu, L. Chen, G. Luo, L. Liu, W. Zhu, P. Zhang, C. Zhang, W. Wu, L. Chen and P. Yan, Study on Synthesis and Biological Evaluation of 3-Aryl Substituted Xanthone Derivatives as Novel and Potent Tyrosinase Inhibitors, *Chemical and Pharmaceutical Bulletin*, 2019, **67**, 1232-1241.
120. E. Yuanita, H. D. Pranowo, M. Mustofa, R. T. Swasono, J. Syahri and J. Jumina, Synthesis, Characterization and Molecular Docking of Chloro-substituted Hydroxyxanthone Derivatives, *Chemistry Journal of Moldova*, 2019, **14**, 68-76.
121. X. D. Kou, L. L. Song, Y. H. Wang, Q. Yu, H. Ju, A. H. Yang and R. Shen, Design, synthesis and anti-Alzheimer's disease activity study of xanthone derivatives based on multi-

REVIEW

Organic Chemistry Frontiers

- target strategy, *Bioorganic & Medicinal Chemistry Letters*, 2020, **30**.
122. S. Genovese, S. Fiorito, M. C. Specchiulli, V. A. Taddeo and F. Epifano, Microwave-assisted synthesis of xanthenes promoted by ytterbium triflate, *Tetrahedron Letters*, 2015, **56**, 847-850.
123. J. Hu, E. A. Adogla, Y. Ju, D. Fan and Q. Wang, Copper-catalyzed ortho-acylation of phenols with aryl aldehydes and its application in one-step preparation of xanthenes, *Chemical Communications*, 2012, **48**, 11256-11258.
124. C. A. Menendez, F. Nador, G. Radivoy and D. C. Gerbino, One-Step Synthesis of Xanthenes Catalyzed by a Highly Efficient Copper-Based Magnetically Recoverable Nanocatalyst, *Organic Letters*, 2014, **16**, 2846-2849.
125. T. Ramani, P. Umadevi, K. L. Prasanth and B. Sreedhar, Synthesis of ortho-Hydroxybenzophenones Catalyzed by Magnetically Retrievable Fe<sub>3</sub>O<sub>4</sub> Nanoparticles under Ligand-Free Conditions, *European Journal of Organic Chemistry*, 2013, **2013**, 6021-6026.
126. A. Venkanna, P. V. K. Goud, P. V. Prasad, M. Shanker and P. V. Rao, One-Step Synthesis of Xanthenes and Thio-Xanthenes by Using Magnetically Separable Copper ferrite Nanoparticles as a Catalyst under Ligand Free Conditions, *Chemistryselect*, 2016, **1**, 2271-2276.
127. H. S. Steingruber, P. Mendioroz, A. S. Diez and D. C. Gerbino, A Green Nanopalladium-Supported Catalyst for the Microwave-Assisted Direct Synthesis of Xanthenes, *Synthesis*, 2020, **52**, 619-628.
128. C. R. Shen and X. F. Wu, Selective Preparation of Xanthenes from 2-Bromofluorobenzenes and Salicylaldehydes via Palladium-Catalyzed Acylation-SNAr Approach, *Synlett*, 2016, **27**, 1269-1273.
129. A. V. Dubrovskiy and R. C. Larock, Intermolecular C-O addition of carboxylic acids to arynes: synthesis of o-hydroxyaryl ketones, xanthenes, 4-chromanones, and flavones, *Tetrahedron*, 2013, **69**, 2789-2798.
130. Y. Fujimoto, H. Yanai and T. Matsumoto, Concise Total Synthesis of Elliptoxanthone A by Utilizing Aromatic Oxy-Cope Rearrangement for Efficient C-Isoprenylation of Xanthone Skeleton, *Synlett*, 2016, **27**, 2229-2232.
131. A. Little and J. A. Porco, Total Syntheses of Graphisin A and Sydownin B, *Organic Letters*, 2012, **14**, 2862-2865.
132. T. Xie, C. Zheng, K. Chen, H. He and S. Gao, Asymmetric Total Synthesis of the Complex Polycyclic Xanthone FD-594, *Angewandte Chemie International Edition*, 2020, DOI: 10.1002/anie.201915787, Ahead of Print.
133. S. Ito, T. Kitamura, S. Arulmozhiraja, K. Manabe, H. Tokiwa and Y. Suzuki, Total Synthesis of Termicalcicolanone A via Organocatalysis and Regioselective Claisen Rearrangement, *Organic Letters*, 2019, **21**, 2777-2781.
134. D. D. Xu, Y. Nie, X. Z. Liang, L. Ji, S. Y. Hu, Q. D. You, F. Wang, H. C. Ye and J. X. Wang, A Concise and Efficient Total Synthesis of alpha-Mangostin and beta-Mangostin from *Garcinia mangostana*, *Natural Product Communications*, 2013, **8**, 1101-1103.
135. X. Wei, D. L. Liang, Q. Wang, X. B. Meng and Z. J. Li, Total synthesis of mangiferin, homomangiferin, and neomangiferin, *Organic & Biomolecular Chemistry*, 2016, **14**, 8821-8831.
136. Q. X. Qin, J. Yang, B. Yang, M. Liu, D. Xiao and X. M. Yang, A facile synthesis of norathyriol, *Journal of Chemical Research*, 2013, DOI: 10.3184/174751912x13547158602399, 38-39.
137. M. Dai, X. Yuan, J. Kang, Z. J. Zhu, R. C. Yue, H. Yuan, B. Y. Chen, W. D. Zhang, R. H. Liu and Q. Y. Sun, Synthesis and biological evaluation of phenyl substituted polyoxygenated xanthone derivatives as anti-hepatoma agents, *European Journal of Medicinal Chemistry*, 2013, **69**, 159-166.
138. K. M. Elbel, G. Guizzunti, M. A. Theodoraki, J. Xu, A. Batova, M. Dakanali and E. A. Theodorakis, A-ring oxygenation modulates the chemistry and bioactivity of caged *Garcinia xanthenes*, *Organic & Biomolecular Chemistry*, 2013, **11**, 3341-3348.
139. C. M. Liu, M. Zhang, Z. H. Zhang, S. B. Zhang, S. M. Yang, A. Zhang, L. J. Yin, S. Swarts, S. Vidyasagar, L. R. Zhang and P. Okunieff, Synthesis and anticancer potential of novel xanthone derivatives with 3,6-substituted chains, *Bioorganic & Medicinal Chemistry*, 2016, **24**, 4263-4271.
140. P. H. Wang, L. L. Jiang, Y. Cao, X. D. Zhang, B. J. Chen, S. Y. Zhang, K. Huang, D. Y. Ye and L. Zhou, Xanthone derivatives as phosphoglycerate mutase 1 inhibitors: Design, synthesis, and biological evaluation, *Bioorganic & Medicinal Chemistry*, 2018, **26**, 1961-1970.
141. M. L. N. Rao and B. S. Ramakrishna, Rh-catalyzed direct synthesis of 2,2'-dihydroxybenzophenones and xanthenes, *Rsc Advances*, 2016, **6**, 75505-75511.
142. X. J. Zhang, S. F. Ye, Y. Zhang, H. Y. Meng, M. Q. Zhang, W. L. Gao and Q. D. You, MICROWAVE-ASSISTED EFFICIENT AND GREEN SYNTHESIS OF HYDROXYXANTHONE IN WATER, *Synthetic Communications*, 2012, **42**, 2952-2958.
143. X. J. Zhang, L. Yang, Y. Wu, J. Y. Du, Y. L. Mao, X. Wang, S. J. Luan, Y. H. Lei, X. Li, H. P. Sun and Q. D. You, Microwave-assisted transition-metal-free intramolecular Ullmann-type O-arylation in water for the synthesis of xanthenes and azaxanthenes, *Tetrahedron Letters*, 2014, **55**, 4883-4887.
144. S. Fuse, K. Matsumura, K. Johmoto, H. Uekusa, H. Tanaka, T. Hirose, T. Sunazuka, S. Omura and T. Takahashi, The Design, Synthesis, and Evaluation of 1,5,7-Trisubstituted-3-Pyridyl-Xanthenes for Use as Insecticides Starting from PirypropeneA, *Chemistry - A European Journal*, 2016, **22**, 18450-18455.
145. M. M. Johnson, J. M. Naidoo, M. A. Fernandes, E. M. Mmutlane, W. A. L. van Otterlo and C. B. de Koning, CAN-Mediated Oxidations for the Synthesis of Xanthenes and Related Products (vol 75, pg 8701, 2010), *Journal of Organic Chemistry*, 2019, **84**, 472-472.
146. J. Dam, M. L. Bode and C. B. de Koning, Ceric Ammonium Sulfate (CAS) Mediated Oxidations of Benzophenones Possessing a Phenolic Substituent for the Synthesis of Xanthenes and Related Products, *Journal of Organic Chemistry*, 2019, **84**, 150-160.
147. N. Marquise, P. Harford, F. Chevallier, T. Roisnel, V. Dorcet, a.-I. Gagez, S. Sablé, L. Picot, V. Thiéry, A. Wheatley, P. Gros and F. Mongin, Synthesis of azafluorenones and related compounds using deprotection-arylation followed by intramolecular direct arylation, *Tetrahedron*, 2013, **69**, 10123.
148. F. Marsais, F. Trécourt, P. Bréant and G. Quéguiner, Directed lithiation of 4-halopyridines: Chemoselectivity, regioselectivity and application to synthesis, *Journal of Heterocyclic Chemistry*, 1988, **25**, 81-87.

149. W. Qian, J. Brown, J. J. Chen and Y. Cheng, Regioselective synthesis of multiply halogenated azaxanthenes, *Tetrahedron Letters*, 2014, **55**, 7229-7232.
150. F. Trécourt, G. Breton, V. Bonnet, F. Mongin, F. Marsais and G. Quéguiner, New Syntheses of Substituted Pyridines via Bromine–Magnesium Exchange, *Tetrahedron*, 2000, **56**, 1349-1360.
151. C. Fernandes, L. Oliveira, M. E. Tiritan, L. Leitao, A. Pozzi, J. B. Noronha-Matos, P. Correia-de-Sa and M. M. Pinto, Synthesis of new chiral xanthone derivatives acting as nerve conduction blockers in the rat sciatic nerve, *European Journal of Medicinal Chemistry*, 2012, **55**, 1-11.
152. W. J. Liu, D. S. Mei and W. H. Duan, Synthesis of 1,7-dimethoxy-2-hydroxyxanthone, a natural product with potential activity on erectile dysfunction, *Chinese Chemical Letters*, 2013, **24**, 515-517.
153. N. Fukushima, N. Ieda, M. Kawaguchi, K. Sasakura, T. Nagano, K. Hanaoka, N. Miyata and H. Nakagawa, Development of photo-controllable hydrogen sulfide donor applicable in live cells, *Bioorganic & Medicinal Chemistry Letters*, 2015, **25**, 175-178.
154. P. Lorenz, A. Heller, M. Bunse, M. Heinrich, M. Berger, J. Conrad, F. C. Stintzing and D. R. Kammerer, Structure Elucidation of the Main Tetrahydroxyxanthenes of Hypericum Seeds and Investigations into the Testa Structure, *Chemistry & Biodiversity*, 2018, **15**.
155. C. Hoppmann, U. Alexiev, E. Irran and K. Ruck-Braun, Synthesis and fluorescence of xanthone amino acids, *Tetrahedron Letters*, 2013, **54**, 4585-4587.
156. M. Presset, J. Paul, G. N. Cherif, N. Ratnam, N. Laloi, E. Leonel, C. Gosmini and E. Le Gall, Co-Catalyzed Barbier Reactions of Aromatic Halides with Aromatic Aldehydes and Imines, *Chemistry – A European Journal*, 2019, **25**, 4491-4495.
157. S. Sarkar and N. Tadigoppula, PhI(OAc)(2)-BF<sub>3</sub>-OEt<sub>2</sub> mediated domino imine activation, intramolecular C-C bond formation and beta-elimination: new approach for the synthesis of fluorenones, xanthenes and phenanthridines, *Rsc Advances*, 2014, **4**, 40964-40968.
158. Z. Z. Shi and F. Glorius, Synthesis of fluorenones via quaternary ammonium salt-promoted intramolecular dehydrogenative arylation of aldehydes, *Chemical Science*, 2013, **4**, 829-833.
159. P. Wang, H. Rao, R. Hua and C.-J. Li, Rhodium-Catalyzed Xanthone Formation from 2-Aryloxybenzaldehydes via Cross-Dehydrogenative Coupling (CDC), *Organic Letters*, 2012, **14**, 902-905.
160. S. Wertz, D. Leifert and A. Studer, Cross Dehydrogenative Coupling via Base-Promoted Homolytic Aromatic Substitution (BHAS): Synthesis of Fluorenones and Xanthenes, *Organic Letters*, 2013, **15**, 928-931.
161. S. K. Manna, S. L. K. Manda and G. Panda, [RuCl<sub>2</sub>(p-cymene)<sub>2</sub>]<sub>2</sub> catalyzed cross dehydrogenative coupling (CDC) toward xanthone and fluorenone analogs through intramolecular C–H bond functionalization reaction, *Tetrahedron Letters*, 2014, **55**, 5759-5763.
162. H. Bao, X. Hu, J. Zhang and Y. Liu, Cu(0)/Selectfluor system-catalyzed intramolecular Csp<sup>2</sup>-H/Csp<sup>2</sup>-H cross-dehydrogenative coupling (CDC), *Tetrahedron*, 2019, **75**, 130533pp.
163. J. Chen and F. R. Sheykhamad, Intramolecular cross-dehydrogenative coupling of benzaldehyde derivatives: A novel and efficient route to benzocyclic ketones, *Journal of the Chinese Chemical Society*, 2019, DOI: 10.1002/jccs.201900214.
164. H. Rao, X. Ma, Q. Liu, Z. Li, S. Cao and C.-J. Li, Metal-Free Oxidative Coupling: Xanthone Formation via Direct Annulation of 2-Aryloxybenzaldehyde using Tetrabutylammonium Bromide as a Promoter in Aqueous Medium, *Advanced Synthesis & Catalysis*, 2013, **355**, 2191-2196.
165. J. Tang, S. J. Zhao, Y. Y. Wei, Z. J. Quan and C. D. Huo, CBr<sub>4</sub> promoted intramolecular aerobic oxidative dehydrogenative arylation of aldehydes: application in the synthesis of xanthenes and fluorenones, *Organic & Biomolecular Chemistry*, 2017, **15**, 1589-1592.
166. H. Jiang, G. Mao, H. Wu, Q. An, M. Zuo, W. Guo, C. Xu, Z. Sun and W. Chu, Synthesis of dibenzocycloketones by acyl radical cyclization from aromatic carboxylic acids using methylene blue as a photocatalyst, *Green Chemistry*, 2019, **21**, 5368-5373.
167. H. F. Motiwala, R. H. Vekariya and J. Aubé, Intramolecular Friedel–Crafts Acylation Reaction Promoted by 1,1,1,3,3,3-Hexafluoro-2-propanol, *Organic Letters*, 2015, **17**, 5484-5487.
168. N. Jiang, S.-Y. Li, S.-S. Xie, H. Yao, H. Sun, X.-B. Wang and L.-Y. Kong, FeCl<sub>3</sub> and ether mediated direct intramolecular acylation of esters and their application in efficient preparation of xanthone and chromone derivatives, *RSC Advances*, 2014, **4**, 63632-63641.
169. S. Kruger and F. G. Mann, Xanthenes and thioxanthenes. Part VI. The preparation and properties of 9-thia-3-aza-anthrone, *Journal of the Chemical Society (Resumed)*, 1955, DOI: 10.1039/JR9550002755, 2755-2763.
170. F. G. Mann and J. H. Turnbull, 164. Xanthenes and thioxanthenes. Part III. The synthesis of 9-oxa-1-aza-anthrone (4-azaxanthone), *Journal of the Chemical Society (Resumed)*, 1951, DOI: 10.1039/JR9510000761, 761-762.
171. J. Li, C. Jin and W. Su, Synthesis of 1-Azaxanthenes and 1-Azathioxanthenes by Yb(OTf)<sub>3</sub>/TfOH Co-Catalyzed Intramolecular Friedel–Crafts Reaction, *Heterocycles*, 2010, **81**, 2555-2567.
172. O. Epstein, M. C. Bryan, A. C. Cheng, K. Derakhchan, T. A. Dineen, D. Hickman, Z. Hua, J. B. Human, C. Kreiman, I. E. Marx, M. M. Weiss, R. C. Wahl, P. H. Wen, D. A. Whittington, S. Wood, X. M. Zheng, R. T. Fremneau, R. D. White and V. F. Patel, Lead Optimization and Modulation of hERG Activity in a Series of Aminooxazoline Xanthene  $\beta$ -Site Amyloid Precursor Protein Cleaving Enzyme (BACE1) Inhibitors, *Journal of Medicinal Chemistry*, 2014, **57**, 9796-9810.
173. H. M. T. Albuquerque, C. M. M. Santos, J. A. S. Cavaleiro and A. M. S. Silva, (E)-2-(4-Arylbut-1-en-3-yn-1-yl)chromones as Synthons for the Synthesis of Xanthone-1,2,3-triazole Dyads, *European Journal of Organic Chemistry*, 2015, **2015**, 4732-4743.
174. H. M. T. Albuquerque, C. M. M. Santos, J. A. S. Cavaleiro and A. M. S. Silva, First intramolecular Diels–Alder reactions using chromone derivatives: synthesis of chromeno 3,4-b xanthenes and 2-(benzo c chromenyl)-chromones, *New Journal of Chemistry*, 2018, **42**, 4251-4260.
175. C. Proenca, H. M. T. Albuquerque, D. Ribeiro, M. Freitas, C. M. M. Santos, A. M. S. Silva and E. Fernandes, Novel chromone and xanthone derivatives: Synthesis and

REVIEW

Organic Chemistry Frontiers

- ROS/RNS scavenging activities, *European Journal of Medicinal Chemistry*, 2016, **115**, 381-392.
176. A. Bornadiego, J. Diaz and C. F. Marcos, Synthesis of 4-Aminoxanthenes by an Uncatalyzed, Multicomponent Reaction, *Advanced Synthesis & Catalysis*, 2014, **356**, 718-722.
177. A. Bornadiego, J. Diaz and C. F. Marcos, Regioselective Tandem 4+1 - 4+2 Synthesis of Amino-Substituted Dihydroxanthenes and Xanthenes, *Journal of Organic Chemistry*, 2015, **80**, 6165-6172.
178. A. Bornadiego, J. Diaz and C. F. Marcos, Tandem synthesis of 4-aminoxanthenes is controlled by a water-assisted tautomerization: A general straightforward reaction, *Organic & Biomolecular Chemistry*, 2019, **17**, 1410-1422.
179. M. Cheng, J. Yan, F. Hu, H. Chen and Y. Hu, Palladium-catalyzed cascade reactions of 3-iodochromones with aryl iodides and norbornadiene leading to annulated xanthenes, *Chemical Science*, 2013, **4**, 526-530.
180. C. Ghosh, D. K. SinhaRoy and K. K. Mukhopadhyay, Heterocyclic systems. Part 6. Reactions of 4-oxo-4H-[1]benzopyran-3-carbonitriles with hydrazine, phenylhydrazine, hydroxylamine, and some reactive methylene compounds, *Journal of the Chemical Society, Perkin Transactions 1*, 1979, DOI: 10.1039/P19790001964, 1964-1968.
181. A. Nohara, T. Ishiguro, K. Ukawa, H. Sugihara, Y. Maki and Y. Sanno, Studies on Antianaphylactic Agents. 7. Synthesis of Antiallergic 5-Oxo-5H-[1]benzopyrano[2,3-b]pyridines, *Journal of Medicinal Chemistry*, 1985, **28**, 559-568.
182. P. Langer and B. Appel, Efficient synthesis of benzopyrano[2,3-b]pyridines by sequential reactions of 1,3-bis-silyl enol ethers with 3-cyanobenzopyrylium triflates, *Tetrahedron Letters*, 2003, **44**, 5133-5135.
183. A. G. P. R. Figueiredo, A. C. Tomé, A. M. S. Silva and J. A. S. Cavaleiro, Reaction of chromone-3-carbaldehyde with  $\alpha$ -amino acids—syntheses of 3- and 4-(2-hydroxybenzoyl)pyrroles, *Tetrahedron*, 2007, **63**, 910-917.
184. N. Poomathi, P. T. Perumal and S. Ramakrishna, An efficient and eco-friendly synthesis of 2-pyridones and functionalized azaxanthone frameworks via indium triflate catalyzed domino reaction, *Green Chemistry*, 2017, **19**, 2524-2529.
185. Q. Yang, R. Wang, J. Han, C. C. Li, T. Wang, Y. Liang and Z. T. Zhang, Photo-induced tandem cyclization of 3-iodoflavones with electron rich five-membered heteroarenes, *Rsc Advances*, 2017, **7**, 43206-43211.
186. Y. Liu, S. Jin, L. Huang and Y. Hu, Phase Transfer Reagent Promoted Tandem Ring-Opening and Ring-Closing Reactions of Unique 3-(1-Alkynyl) Chromones, *Organic Letters*, 2015, **17**, 2134-2137.
187. S. A. French, M. R. Clark, R. J. Smith, T. Brind and B. C. Hawkins, Expedient synthesis of xanthenes and multifunctionalized chromones from 1,1-diacyl cyclopropanes, *Tetrahedron*, 2018, **74**, 5340-5350.
188. K. Goto, M. Terasawa and Y. Maruyama, Anti-Allergic Activities of a New Benzopyranopyridine Derivative Y-12,141 in Rats, *International Archives of Allergy and Immunology*, 1979, **59**, 13-19.
189. P. G. Dormer, K. K. Eng, R. N. Farr, G. R. Humphrey, J. C. McWilliams, P. J. Reider, J. W. Sager and R. P. Volante, Highly regioselective Friedlander annulations with unmodified ketones employing novel amine catalysts: syntheses of 2-substituted quinolines, 1,8-naphthyridines, and related heterocycles, *Journal of Organic Chemistry*, 2003, **68**, 467-477.
190. H. Waldmann, G. V. Karunakar and K. Kumar, Gold(III)-Mediated Aldol Condensations Provide Efficient Access to Nitrogen Heterocycles, *Organic Letters*, 2008, **10**, 2159-2162.
191. B. Duda, S. N. Tverdomed and G. V. Roschenthaler, CF<sub>2</sub>-containing acetylenephosphonates in heterocyclization reactions: the first synthesis of 2-difluoromethyl azaxanth-3-ylphosphonates, *Organic & Biomolecular Chemistry*, 2011, **9**, 8228-8232.
192. K. Otrubova, A. E. Fitzgerald and N. S. Mani, A novel entry to xanthenes by an intramolecular Diels-Alder reaction involving 2-(1,2-dichlorovinyl) aryl dienones, *Tetrahedron*, 2018, **74**, 5715-5724.
193. Y. He, S. H. Guo, X. S. Fan, C. H. Guo and X. Y. Zhang, One-pot synthesis of 12H-benzo b xanthen-12-ones and naphtha 2,3-b furans from non-naphthalene substrates, *Tetrahedron Letters*, 2014, **55**, 4747-4752.
194. Z. L. Xiong, X. H. Zhang, Y. M. Li, X. S. Peng, J. Y. Fu, J. J. Guo, F. K. Xie, C. G. Jiang, B. Lin, Y. X. Liu and M. S. Cheng, Syntheses of 12H-benzo a xanthen-12-ones and benzo a acridin-12(7H)-ones through Au(I)-catalyzed Michael addition/6-endo-trig cyclization/aromatization cascade annulation, *Organic & Biomolecular Chemistry*, 2018, **16**, 7361-7374.
195. G. X. Z. Liu, C. Wu, B. F. Chen, R. He and C. Chen, Concise synthesis of xanthenes by the tandem etherification-Acylation of diaryliodonium salts with salicylates, *Chinese Chemical Letters*, 2018, **29**, 985-988.
196. X. C. Cheng, Y. Y. Zhou, F. F. Zhang, K. Zhu, Y. Y. Liu and Y. Z. Li, Base-Promoted Tandem Reaction Involving Insertion into Carbon-Carbon sigma-Bonds: Synthesis of Xanthone and Chromone Derivatives, *Chemistry - A European Journal*, 2016, **22**, 12655-12659.
197. Y. Fujimoto, R. Itakura, H. Hoshi, H. Yanai, Y. Ando, K. Suzuki and T. Matsumoto, Novel One-Pot Synthesis of Xanthenes via Sequential Fluoride Ion-Promoted Fries-Type Rearrangement and Nucleophilic Aromatic Substitution, *Synlett*, 2013, **24**, 2575-2580.
198. J. Kristensen, M. Begtrup and P. Vedsø, ChemInform Abstract: Preparation of 5-Acyl- and 5-Aryl-Substituted 1-(Benzyloxy)pyrazoles via Directed Ortho-Lithiation/Transmetalation and Palladium Catalyzed Cross-Coupling, *Synthesis*, 1998, **30**, 1604-1608.
199. J. Felding, P. Uhlmann, J. Kristensen, P. Vedsø and M. Begtrup, Palladium-Catalyzed Arylation and Acylation of 1-Benzyloxy-1,2,3-triazole Through a Directed Ortho Lithiation-Transmetalation Strategy, *Synthesis*, 1998, **1998**, 1181-1184.
200. J. L. Kristensen, P. Vedsø and M. Begtrup, Synthesis of novel azaxanthenes derived from N-hydroxyazoles, *Tetrahedron*, 2002, **58**, 2397-2404.
201. G. Singh, R. Singh, N. K. Girdhar and M. P. S. Ishar, A versatile route to 2-alkyl-/aryl-amino-3-formyl- and hetero-annulated-chromones, through a facile nucleophilic substitution at C2 in 2-(N-methylanilino)-3-formylchromones, *Tetrahedron*, 2002, **58**, 2471-2480.
202. F. Mann and J. Reid, 382. Xanthenes and thioxanthenes. Part IV. The preparation and properties of 9-oxa-1-aza-anthrone and 9-thia-1-aza-anthrone, *Journal of The*



- Chemical Society (resumed)*, 1952, DOI: 10.1039/jr9520002057.
203. G. Kolokythas, N. Pouli, P. Marakos, H. Pratsinis and D. Kletsas, Design, synthesis and antiproliferative activity of some new azapyranoxanthenone aminoderivatives, *European Journal of Medicinal Chemistry*, 2006, **41**, 71-79.
  204. D. E. Daia, C. D. Gabbutt, B. M. Heron, J. D. Hepworth, M. B. Hursthouse and K. M. A. Malik, Conjugate additions of lithium dialkynylcuprates [(RC)C<sub>2</sub>CuLi] to activated chromones. Unexpected formation of the 6H-bis[1]benzopyrano[2,3-b:3',4'-e]pyridine system, *Tetrahedron Letters*, 2003, **44**, 1461-1464.
  205. Z. N. Siddiqui, One pot synthesis of new benzopyranopyridines via Friedlander condensation, *Tetrahedron Letters*, 2012, **53**, 4974-4978.
  206. P. Biswas, J. Ghosh, T. Sarkar, D. Jana and C. Bandyopadhyay, Design and synthesis of dichromeno[2,3-b:3',2'-e]pyridine-12,14-dione to evaluate its optical properties, *New Journal of Chemistry*, 2017, **41**, 3887-3893.
  207. C. K. Ghosh, K. Bhattacharya and C. Ghosh, Benzopyrans - XXXIII. [4+2]Cycloaddition of N,N-dimethylhydrazones and anils of 2-unsubstituted and 2-methyl-4-oxo-4H-1-benzopyran-3-carboxaldehyde with N-phenylmaleimide, *Tetrahedron*, 1994, **50**, 4905-4912.
  208. T. U. Kumar, D. Roy and A. Bhattacharya, Iron(III) catalyzed direct C-H functionalization at the C-3 position of chromone for the synthesis of fused chromeno-quinoline scaffolds, *Tetrahedron Letters*, 2019, **60**, 1895-1898.
  209. T. Schurreit, 4-Hydroxy-2H-[1]benzopyran-2-on als Baustein zur Synthese von Bisbenzopyranopyridinen, *Archiv der Pharmazie*, 1987, **320**, 500-506.
  210. Y. Panneerselvam and B. s. R. Reddy, One pot synthesis of 1-azaxanthenes via Friedlander condensation, *Indian Journal of Chemistry Section B*, 2016, **55**, 83-87.
  211. C. P. Montgomery, D. Parker and L. Lamarque, Effective and efficient sensitisation of terbium luminescence at 355 nm with cell permeable pyrazoyl-1-azaxanthe macrocyclic complexes, *Chemical Communications*, 2007, DOI: 10.1039/B709805G, 3841-3843.
  212. R. Martín, L. B. Jiménez, M. Álvaro, J. C. Scaiano and H. García, Photoinduced Formation and Characterization of Electron-Hole Pairs in Azaxanthylum-Derivatized Short Single-Walled Carbon Nanotubes, *Chemistry – A European Journal*, 2009, **15**, 8751-8759.
  213. H.-J. Cho, M.-J. Jung, Y. Kwon and Y. Na, Oxiranylmethoxy or thiiranylmethoxy-azaxanthenes and -acridone analogues as potential topoisomerase I inhibitors, *Bioorganic & Medicinal Chemistry Letters*, 2009, **19**, 6766-6769.
  214. T. Ghosh and C. Bandyopadhyay, Reactions of 2-Amino-4H-1-benzopyran-4-one with 4-Oxo-4H-1-benzopyran-3-carbaldehydes, *Journal of Heterocyclic Chemistry - J HETEROCYCL CHEM*, 2006, **43**, 1431-1434.
  215. D. Imperio, G. B. Giovenzana, G.-I. Law, D. Parker and J. W. Walton, Synthesis and comparative anion binding profiles of two di-aqua Eu(III) complexes, *Dalton Transactions*, 2010, **39**, 9897-9903.
  216. G.-L. Law, R. Pal, L. O. Palsson, D. Parker and K.-L. Wong, Responsive and reactive terbium complexes with an azaxanthe sensitizer and one naphthyl group: applications in ratiometric oxygen sensing in vitro and in regioselective cell killing, *Chemical Communications*, 2009, DOI: 10.1039/B920222F, 7321-7323.
  217. H. Nouri, C. Cadiou, L. M. Lawson-Daku, A. Hauser, S. Chevreux, I. Dechamps-Olivier, F. Lachaud, R. Ternane, M. Trabelsi-Ayadi, F. Chuburu and G. Lemerrier, A modified cyclen azaxanthe ligand as a new fluorescent probe for Zn<sup>2+</sup>, *Dalton Trans*, 2013, **42**, 12157-12164.
  218. A. Rampa, A. Bisi, P. Valenti, M. Recanatini, A. Cavalli, V. Andrisano, V. Cavrini, L. Fin, A. Buriani and P. Giusti, Acetylcholinesterase Inhibitors: Synthesis and Structure-Activity Relationships of ω-[N-Methyl-N-(3-alkylcarbamoyloxyphenyl)-methyl]aminoalkoxyheteroaryl Derivatives, *Journal of Medicinal Chemistry*, 1998, **41**, 3976-3986.
  219. S. Gobbi, Q. Z. Hu, M. Negri, C. Zimmer, F. Belluti, A. Rampa, R. W. Hartmann and A. Bisi, Modulation of Cytochromes P450 with Xanthone-Based Molecules: From Aromatase to Aldosterone Synthase and Steroid 11 beta-Hydroxylase Inhibition, *Journal of Medicinal Chemistry*, 2013, **56**, 1723-1729.
  220. N. Szkaradek, A. Rapacz, K. Pytka, B. Filipek, D. Zelazczyk, P. Szafranski, K. Sloczynska and H. Marona, Cardiovascular activity of the chiral xanthone derivatives, *Bioorganic & Medicinal Chemistry*, 2015, **23**, 6714-6724.
  221. N. Szkaradek, D. Sypniewski, A. M. Waszkielewicz, A. Gunia-Krzyzak, A. Galilejczyk, S. Galka, H. Marona and I. Bednarek, Synthesis and in vitro Evaluation of the Anticancer Potential of New Aminoalkanol Derivatives of Xanthone, *Anti-Cancer Agents in Medicinal Chemistry*, 2016, **16**, 1587-1604.
  222. M. Kubacka, N. Szkaradek, S. Mogilski, K. Panczyk, A. Siwek, A. Grybos, B. Filipek, P. Zmudzki, H. Marona and A. M. Waszkielewicz, Design, synthesis and cardiovascular evaluation of some aminoisopropanol derivatives of xanthone, *Bioorganic & Medicinal Chemistry*, 2018, **26**, 3773-3784.
  223. N. Szkaradek, A. Rapacz, K. Pytka, B. Filipek, A. Siwek, M. Cegla and H. Marona, Synthesis and preliminary evaluation of pharmacological properties of some piperazine derivatives of xanthone, *Bioorganic & Medicinal Chemistry*, 2013, **21**, 514-522.
  224. M. Franceschin, D. Nocioni, A. Biroccio, E. Micheli, S. Cacchione, C. Cingolani, A. Venditti, P. Zizza, A. Bianco and A. Altieri, Design and synthesis of a new dimeric xanthone derivative: enhancement of G-quadruplex selectivity and telomere damage, *Organic & Biomolecular Chemistry*, 2014, **12**, 9572-9582.
  225. D. Zelazczyk, M. Jakubczyk, K. Pytka, A. Rapacz, M. Walczak, P. Janiszewska, K. Panczyk, H. Marona, A. M. Waszkielewicz, P. Zmudzki and K. Sloczynska, Design, synthesis and evaluation of activity and pharmacokinetic profile of new derivatives of xanthone and piperazine in the central nervous system, *Bioorganic & Medicinal Chemistry Letters*, 2019, **29**, 126679.
  226. S. Motavallizadeh, A. Fallah-Tafti, S. Maleki, A. N. Shirazi, M. Pordeli, M. Safavi, S. K. Ardestani, S. Asd, R. Tiwari, D. Oh, A. Shafiee, A. Foroumadi, K. Parang and T. Akbarzadeh, Synthesis and evaluation of antiproliferative activity of substituted N-(9-oxo-9H-xanthen-4-yl)benzenesulfonamides, *Tetrahedron Letters*, 2014, **55**, 373-375.



REVIEW

Organic Chemistry Frontiers

227. X. Fei, M. Jo, B. Lee, S. B. Han, K. Lee, J. K. Jung, S. Y. Seo and Y. S. Kwak, Synthesis of xanthone derivatives based on alpha-mangostin and their biological evaluation for anti-cancer agents, *Bioorganic & Medicinal Chemistry Letters*, 2014, **24**, 2062-2065.
228. D. Giallombardo, A. C. Nevin, W. Lewis, C. C. Nawrat, R. R. A. Kitson and C. J. Moody, Synthesis of toxyloxanthone B, *Tetrahedron*, 2014, **70**, 1283-1288.
229. T. S. Ehianeta, S. Laval and B. Yu, Bio- and chemical syntheses of mangiferin and congeners, *Biofactors*, 2016, **42**, 445-458.
230. X. Wei, D. L. Liang, M. H. Ning, Q. Wang, X. B. Meng and Z. J. Li, Semi-synthesis of neomangiferin from mangiferin, *Tetrahedron Letters*, 2014, **55**, 3083-3086.
231. Y. Fujimoto, Y. Watabe, H. Yanai, T. Taguchi and T. Matsumoto, An Efficient Isoprenylation of Xanthenes at the C1 Position by Utilizing Anion-Accelerated Aromatic Oxy-Cope Rearrangement, *Synlett*, 2016, **27**, 848-853.
232. O. Talhi and A. M. S. Silva, Organic Synthesis of C-Prenylated Phenolic Compounds, *Current Organic Chemistry*, 2013, **17**, 1067-1102.
233. R. S. Wang, R. D. Chen, J. H. Li, X. Liu, K. B. Xie, D. W. Chen, Y. Peng and J. G. Dai, Regiospecific Prenylation of Hydroxyxanthenes by a Plant Flavonoid Prenyltransferase, *Journal of Natural Products*, 2016, **79**, 2143-2147.
234. X. Li, Y. Wu, Y. Y. Wang, Q. D. You and X. J. Zhang, "Click Chemistry" Synthesis of Novel Natural Product-Like Caged Xanthenes Bearing a 1,2,3-Triazole Moiety with Improved Druglike Properties as Orally Active Antitumor Agents, *Molecules*, 2017, **22**.
235. G. A. Kraus, T. Guney, A. Kempema, J. M. Hyman and B. Parvin, Efficient synthesis of fluorescent rosamines: multifunctional platforms for cellular imaging, *Tetrahedron Letters*, 2014, **55**, 1549-1551.
236. O. A. Akrawi, H. H. Mohammed, T. Patonay, A. Villinger and P. Langer, Synthesis of arylated xanthenes by site-selective Suzuki-Miyaura reactions of the bis(triflate) of 1,3-dihydroxyxanthone, *Tetrahedron*, 2012, **68**, 6298-6304.
237. M. M. Lakouraj, G. Rahpaima and S. M. Mohseni, Synthesis, characterization, and biological activities of organosoluble and thermally stable xanthone-based polyamides, *Journal of Materials Science*, 2013, **48**, 2520-2529.
238. V. Hasantabar, M. M. Lakouraj, E. N. Zare and M. Mohseni, Synthesis, Characterization, and Biological Properties of Novel Bioactive Poly(xanthoneamide-triazole-ethersulfone) and Its Multifunctional Nanocomposite with Polyaniline, *Advances in Polymer Technology*, 2017, **36**.
239. R. Shen, Y. Chen, Z. Li, H. Qi and Y. Wang, Synthesis and biological evaluation of disubstituted amidoxanthenes as potential telomeric G-quadruplex DNA-binding and apoptosis-inducing agents, *Bioorganic & Medicinal Chemistry*, 2016, **24**, 619-626.
240. D. I. S. P. Resende, P. Pereira-Terra, J. Moreira, J. Freitas-Silva, A. Lemos, L. Gales, E. Pinto, E. Sousa, P. M. d. Costa and M. M. M. Pinto, Synthesis of a Small Library of Nature-Inspired Xanthenes and Study of their Antimicrobial Activity, *Molecules*, 2020, **25**, 2405.
241. A. M. Waszkielewicz, A. Gunia, N. Szkaradek, K. Pytko, A. Siwek, G. Satala, A. J. Bojarski, E. Szneler and H. Marona, Synthesis and evaluation of pharmacological properties of some new xanthone derivatives with piperazine moiety, *Bioorganic & Medicinal Chemistry Letters*, 2013, **23**, 4419-4423.

## 1. Proposed title

Recent advances in the synthesis of xanthenes and azaxanthenes

## 2. Authors and affiliations (including co-authors if known)

Diana I. S. P. Resende, <sup>a,b,†</sup> Fernando Durães, <sup>a,b,†</sup> Miguel Maia, <sup>a,b,†</sup> Emília Sousa <sup>a,b,\*</sup> and Madalena M. M. Pinto <sup>a,b</sup>

*CIIMAR - Centro Interdisciplinar de Investigação Marinha e Ambiental, Terminal de Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal.*

*Laboratório de Química Orgânica e Farmacêutica, Faculdade de Farmácia, Universidade do Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal*

† These authors contributed equally to this work.

\*Corresponding author. E-mail address: esousa@ff.up.pt.

## 3. Proposed submission date

3<sup>rd</sup> June 2020

## 4. A brief description of the review and topic to be covered, including:

### a. The current importance of the field

Over the past few decades, strategic advances have been made in medicinal chemistry and organic synthesis that accelerated drug discovery. Diversity-oriented synthesis, photo-induced reactions, biorthogonal chemistry with new and eco-friendly reagents, late-stage diversification, and artificial intelligence are among the approaches involved in some of the recent breakthroughs on drug discovery. More than 85% of all biologically-active chemical entities contain a heterocycle which provides a useful tool for lead optimization. This current review aims to emphasize a particular class of heterocycles, the xanthenes, the use of these advances that allow to speeding up their synthesis and leading optimization process.

Xanthenes are important O-heteroaromatic tricyclic molecules exhibiting a wide range of bioactivities and can represent a showcase to highlight recent techniques and strategies in organic synthesis towards lead discovery and optimization. Several strategies for their synthesis have been developed in recent years, bringing innovation and value to traditional employed methodologies.

### b. What the implications are for the wider scientific community

This review will highlight that the xanthone scaffold still gathers a lot of attention not only from a synthetic point of view but due to the diversity of bioactive derivatives being disclosed, the case-studies of this study. New biological applications are being proposed for xanthenes beyond their pharmacological properties with relevance for chemical biology as sensitizing chromophores. Therefore, it is useful to know the vast array of approaches being used, as they can become a valuable toolbox in the search of novel bioactive heterocycles, emphasizing the importance of this privileged scaffold. Moreover, the site-selective late-stage functionalization approaches highlighted can guide medicinal chemists in the lead optimization not only of (aza)xanthenes but also of isosters like thioxanthenes, acridones, cromones, flavones, among others. This review can help guide medicinal chemists working with heterocycles by providing chemical tools and reaction conditions to both target and diversity-oriented synthesis approaches.

**c. Comments on any other reviews published on a similar topic, justifying why there is room for another review**

The first methods for the synthesis of xanthenes were introduced in by Michael and Kostanecki in 1883-1891 and in the 20th century and until 2012 several extensive reviews have reported the state of the art regarding the synthesis of xanthenes. More recently reviews have focused on the synthesis of a very specific class of derivatives like carboxyxanthenes, hydroxanthenes, arylxanthenes, and thioxanthenes and, although azaxanthenes have demonstrated therapeutic potential and interest as sensitizing chromophores, their reports in medicinal chemistry are still sparse in literature. Therefore, in this review the recent exploitation of both scaffolds was considered from a medicinal chemistry point of view with advanced synthetic methodologies applied for obtaining bioactive derivatives being displayed in the corresponding sections.

**5. Key literatures that you intend to review in your paper**

This review will give an insight of the recent literature disclosed to obtain the xanthone core, exploiting the optimization of well-known procedures as well as disruptive developed methodologies. Their advantages and drawbacks will be discussed, with a focus on their efficiency, eco-friendly conditions and substituents tolerability. Particularly, advanced conditions in cyclization such as metal-free variants of the intramolecular Friedel-Crafts acylation or rhodium-catalyzed cross dehydrogenative coupling of diaryl ethers have been reported. Reaction conditions are following new trends with N-heterocyclic carbenes, quaternary ammonium salts, and photoredox conditions. The syntheses of drug-like azaxanthenes, isosteres of xanthenes, are also explored. Late-stage functionalization of (azo)xanthone moiety is highlighted as a strength in the lead optimization process. With this work, it is expected to provide a useful chemical toolbox to the synthetic medicinal chemistry focusing in these privileged structures.